

SAN DIEGO POLICE DEPARTMENT CRIME LABORATORY





Approved by: Chelsea Carter, Supervising Criminalist November 25, 2019

1.0 INTRODUCTION

1.1 GENERAL GUIDELINES

1.1.1. This manual covers policies, methods, and procedures utilized in the analysis of the typical drugs of abuse that comprise the majority of this section's seized drug casework. It is not meant to cover rare and infrequent submissions. In the case of a non-routine submission, the Criminalist, using sound scientific principles, will select an examination scheme comprised of the types of analyses available to them and outlined in this manual. Standards, controls, and reference materials must be fully documented in the case packet, if applicable. In the event that new procedures, methodology, or instrumentation must be utilized in an analysis, the new method must be validated and approved in accordance with laboratory procedures prior to use.

1.2 UNIT DESCRUTION

- 1.2.1. The Forensic Chemistry Unit is budgeted for eight positions: one Supervising Criminalist, six Criminalists, and one laboratory technician.
- 1.2.2. The unit is located at Ponce Headquarters. Seized drug analysis is performed on the 6th floor in the Porensic Chemistry Unit, located in rooms 617 and 618.
- 1.2.3. The criminalist positions in the unit are governed by civil service requirements that call for a four-year science degree as a minimum expectation.

1.3 UNIT FUNCTIONS

- 1.3.1. This unit performs controlled substance analysis and alcohol analysis.
- 1.3.2. General duties performed include:
 - 1.3.2.1. Performing analysis on suspected controlled substances the form of solids, liquids, pills, plant material, and mushrooms.
 - 1.3.2.2. Court testimony regarding all aspects of analysis and interpretation of results.
- 1.3.3. Combinations of methods are used to identify controlled substances. These methods include:
 - 1.3.3.1. Color tests
 - 1.3.3.2. Microcrystalline tests

S.

- 1.3.3.3. Microscopic examinations
- 1.3.3.4. Raman spectroscopy
- 1.3.3.5. Infrared spectroscopy
- 1.3.3.6. Gas chromatography/mass spectroscopy
- 1.3.3.7. Literature, CD-Rom, and Internet references for pharmaceutical pill identification. TORLABORATORY USE

2.0 PERSONNEL AND JOB DESCRIPTIONS

SUPERVISING CRIMINALIST 2.1

2.1.1.1.

2

2.1.1.3

The duties of the supervisor in the Forensic Chemistry Unit include:

- Supervising the analysis of controlled substances.
 - Ensuring proper procedures are followed.
- Reviewing case packets to ensure proper documentation of analytical procedures.
- nsuring adequate unit staffing levels every day. 2.1.1.4
- ing that new criminalists receive the proper training and 2.1.1.5. En pass appropriate competency tests, written tests, and mock court testimonv.
- Serving as a linison between the contractors, department, 2.1.1.6. district attorney's office, city attorney's office, and other end users of the laporatory.
- Ensuring unit policies are being followed though documentation 2.1.1.7. in logs and records.
- Evaluating employee performa-2.1.1.8.
- 2.1.1.9. Preparing staff reports:

2.1.1.9.1.	Budget requests
2.1.1.9.2.	Monthly unit statistics
2.1.1.9.3.	Special projects

- Acting as an advocate for the staff to upper manage 2.1.1.10. C Ser
- Monitoring and approving electronic time cards. 2.1.1.11.

2.2 CRIMINALIST I & CRIMINALIST II

2.1.4.

2.2.1.5

- 2.2.1. The duties of the Criminalists in the Forensic Chemistry Unit include:
 - 2.2.1.1. Analyzing impounded evidence for controlled substances.
 - 2.2.1.2. Identifying the relevant items to be examined.
 - 2.2.1.3. Providing and documenting correct maintenance of tools, equipment, and instrumentation as needed, and monitoring instruments and arranging for repair as needed.
 - Preparing reagents as needed.

Preparing legible notes and reports.

- 2.2.2.6. Ensuring proper sealing and disposition of evidence, and maintaining proper chain of custody.
- 2.2.1.7. **Performing quarterly checks of reagents and instruments**.
- 2.2.1.8. Keeping the supervisor informed of operations, problems, and unusuar circumstances.
- 2.2.1.9. Maintaining proper public relations.
- 2.2.1.10. Carrying out special projects as requested by supervisor.
- 2.2.1.11. Acting as a technical resource for the Department and others and giving appropriate explanations of conclusions to officers and attorneys in a timely namer.
- 2.2.1.12. Assisting other criminalists with training in analytical and administrative procedures and technical problems.
- 2.2.1.13. Participating in the development of new procedures and validating new instrumentation and equipment as needed.
- 2.2.1.14. Distributing reports to district and city attorn ys when necessary.
- 2.2.1.15. Testifying as a controlled substance analyst expert in court
- 2.2.1.16. Preparing monthly statistics.
- 2.2.1.17. Following laboratory safety procedures.
- 2.2.1.18. Participating in the proficiency test program.

2.3 CRIMINALIST III (Technical Lead)

2.3.1.3.

.3.1.4

- 2.3.1. The duties of the Criminalist III in the Forensic Chemistry Unit include:
 - 2.3.1.1. Coordinating the technical operation of the unit in accordance with quality assurance standards.
 - 2.3.1.2. Creating, reviewing, and/or revising technical policies and procedures prior to final approval by the Quality Manager.

Coordinating training of new and current employees in analytical procedures, and ensuring completion of training documentation.

- Developing technical training plans in the unit.
- 2.3.15. Coordinating, reviewing, and approving new method and instrument validations and verifications, including locumentation, prior to final approval by the Quality Manager.
- 2.3.1.6. Acting as technical reference for the unit Supervisor, Quality Manager and any auditors.
- 2.3.1.7. Acting as a mediator, when necessary, in the technical review process.
- 2.3.1.8. Providing technical consultation, as needed, to members of the unit.
- 2.3.1.9. Working with the supervisor and Quality Manager to ensure criminalist compliance with QA, laboratory, and unit policies and procedures.
- 2.3.1.10. Evaluating new technologies to determine if appropriate for the unit.
- 2.3.1.11. When technical problems are identified, coordinating with the Quality Manager and Supervisor to ensure appropriate corrective actions are taken.
- 2.3.1.12. The duties of the Criminalist III in the Forensic Chemistry Unit also include those of the Criminalist I and II in section 7.2.

2.4 LABORATORY TECHNICIAN

2.4.1.4.

2.4.1.5

- 2.4.1. The duties of the laboratory technician in the Forensic Chemistry Unit include:
 - 2.4.1.1. Ordering, picking up, and stocking supplies for the Unit and coordinating with vendors.
 - 2.4.1.2. Preparing and stocking reagents as needed.
 - 2.4.1.3. Washing lab-ware and filling printers as needed.
 - Assisting in monitoring instruments and arranging for repairs and service as needed.
 - Following proper safety procedures and assisting in safety projects.
 - 2.4.1.6. Keeping the supervisor informed of operations, problems, and inusual circumstances.
 - 2.4.1.7. Miraining proper public relations.
 - 2.4.1.8. Carrying out special projects as requested by the supervisor.
 - 2.4.1.9. Participating in the development of new procedures and validations, is record.
 - 2.4.2.10. Helping maintain the various standards, reagents, and logs, and performing quarterly checks.
 - 2.4.2.11. Maintaining and updating metry Data Sheets.
 - 2.4.2.12. Cleaning common areas in the Vit.
 - 2.4.2.13. Maintaining the chemical inventory for the Unit.
 - 2.4.2.14. Testifying in court if called to do so.
 - 2.4.2.15. Providing assistance to other units of the laboratory when requested and available.
 - 2.4.2.16. Disposing of hazardous material and chemical waste as needed.

3.0 SUBMISSIONS AND HANDLING

3.1 IMPOUND SUBMISSIONS

3.1.1. Seized drug evidence is submitted under an incident number. The items contained within the impound will be identified with unique barcode numbers. A barcode number can be used to identify one item, or multiple items within an impound.

Forensic Chemistry Criminalists receive impounds from the Narcotics Vault or Property personnel.

3.2 ITEMS NOT EXAMINED

3.2.1. The following i empare not routinely analyzed because either the District Attorney's and/or City Attorney's offices do not file on these cases or not enough material is present for analysis. Requests for analysis will be considered on a case oy-case basis by the unit supervisor.

3.2.1.1.	Syringes
3.2.1.2.	Drug paraphenalm
3.2.1.3.	Residue quantities and quantities weighing less than 0.04 grams
3.2.1.	
3.2.1.4.	Marijuana and Marijuana products, including concentrates and edibles
3.2.1.5.	Impounds without subject's name or identifier (except "buy" cases)

3.2.1.6. Liquid other than suspected PCP or GHB-type compounds of less than approximately 0.5ml. Suspected PCP or less than 0.1ml

- 3.2.1.7. Precursors and breakdown products of controlled substances unless they are the only chemicals present for that suspect
- 3.2.1.8. Food items
- 3.2.1.9. Suspected LSD of less than 1 square of paper or gel
- 3.2.1.10. Partial tablets or tablet fragments
- 3.2.1.11. Prescription medications in the subject's name

3.2.2. Items that pose a safety hazard to lab personnel such as: suspected terrorist powders, possible explosives, and materials that react with strong acids and bases used in chemical testing, will not be analyzed.

3.3 SAMPLING PLAN

3.3.1.1.

- 3.3.1. Sampling Plan: The number of items analyzed per defendant is dependent on the charges filed in the case, such as possession or sales related charges.
 - The casework approach for each criminalist with respect to items tested will be to analyze enough items of evidence to meet the charges. For possession cases the Criminalist will test one drug type. There is not a minimum number of items for examination.
 - 3.3.1.2. In general for sales cases, 60% of each drug type will be inalyzed per suspect. Each Criminalist will keep in mind that enhancements exist for possession of certain weights of cocaine and methamphetamine of 1 ounce, 2 ounces, kilos, and so on, ard 1/2 ownce increments for heroin that may require more analysis.
 - 3.3.1.3. The number of items actually tested will be up to the Criminalist and will be r flected in the seized drug report along with the analytical results and weights for those items. No assumptions will be made in the notes or final written product as to the contents of any un ested submissions.

3.4 PRELIMINARY TESTING

- 3.4.1. Cases impounded at Headquarters on Fridays offer the morning pick-up will be given the highest priority on the following Monday morning if no other uncompleted cases are due that day.
- 3.4.2. Headquarter cases impounded on the weekend should be analyzed by the end of business day on the following Monday.
- 3.4.3. Preliminary testing should routinely be completed by the end of the business day on the day the items were received by the Criminalist.
- 3.4.4. Turn-around times are subject to change based on staffing, numbers or examinations required for each impound, and complexity of the analysis required. Complex cases requiring GC/MS analysis are normally completed within three days of submission.
- 3.4.5. The supervisor may reallocate resources and may notify the District Attorney's Office or City Attorney's Office of any delays.
- 3.4.6. If needed, a criminalist may return unanalyzed cases to the Narcotics Vault.

3.5 FINAL TESTING

3.5.1. Receipt of a court subpoena for a seized drug case is the notice that the evidence must be confirmed for trial unless we are notified confirmation is no longer needed. The Criminalist who did the preliminary test will analyze these cases, if possible. It is the Criminalist's responsibility to ensure that the final work is completed and the review process has been completed prior to the court date. Generally, a minimum of five-work days' notice is required for final analysis.

Laboratory requests may be received from other units, such as Homicide or Sex Crimes, for analyses of substances seized during an investigation. The unasupervisor will assign these cases to criminalists. These cases will be analyzed and confirmed, producing a final report.

3.6 BUY PROGPAMS

- 3.6.1. There are two types of buy programs: Buy-Walk and Buy-Bust. Buy-Walk operations involve one purchase of controlled substances with an arrest occurring at a later time. A Buy-Bust operation involves the purchase of controlled substances and an immediate arrest. Those arrestees will be incustody pending arraignment.
- 3.6.2. The narcotics detective should notify the vault and the forensic chemistry supervisor in advance of a by p ogram. The detective will provide the vault staff or the forensic chemistry supervisor with the following information:
 - 3.6.2.1. Name and phone number of primary contact.
 - 3.6.2.2. Operation code name that vill connotated on every impound.
 - 3.6.2.3. Approximate start date and length of program.
 - 3.6.2.4. Approximate number of impounds an icipated.
- 3.6.3. One criminalist will be assigned to work all impounds under a program. Buy program impounds are worked in addition to routine casework.
- 3.6.4. It is the responsibility of the assigned criminalist to ensure that all impounds submitted have been finalized.
- 3.6.5. Criminalists must keep the supervisor apprised of the status of each program
- 3.6.6. Buy program impounds are to be worked as time permits between the higher priority casework.

3.7 IMPOUND RECEIPT AND RETURN

3.7.1. Impounds are generally stored in the Narcotics Vault. Each criminalist must sign for custody of the item from Property or Narcotics Vault personnel.

- 3.7.1.1. When possible, completed cases should be returned to the Vault at the end of each day.
- 3.7.1.2. Daily impound cases not returned to the Vault at the end of the day, will be stored, closed, in locked cabinets in the Forensic Chemistry Unit.

3.7.2. No criminalist should accept an impound directly from an officer. It must be received through the Vault.

3.8 SLALING OF EVIDENCE

- 3.8.1. All evidence shall be sealed prior to being accepted for analysis.
 - 3.8 11. In cases where it is not sealed, the condition will be documented in the notes and the Criminalist will seal the item.
- 3.8.2. Whenever possible, impounds should be opened in a manner that maintains the original evid evid evid and identifying information.
- 3.8.3. No more than one care will be opened at a time for examination.
- 3.8.4. Items analyzed will be the electronic description of the second s
 - 3.8.4.1. Barcode # 3.8.4.2. Initials
- 3.8.5. All analyzed items will be sealed in appropriate packaging.
- 3.8.6. Following examination, the impound will be scared using the department issued evidence tape. The seal should run across the outer opening of the impound. The Criminalist must initial and date across the seal and onto the impound itself.
- 3.8.7. The evidence seal should be placed, whenever possible, in such a location as to not cover another person's seal or any identifying information.

3.9 HANDLING AND REPACKING ITEMS

- 3.9.1. Suspected PCP and Fentanyl cases along with other possible hazardous samples must be handled with gloves and be kept in the hood during analyss. Eye protection is recommended. For Fentanyl cases, a respirator is also recommended along with working the case in the special ductless fume hood located in the lab.
- 3.9.2. Analyzed syringes will not be recapped by criminalists. After removing the cap and dispensing a portion of the contents for analysis, the uncapped syringe will be placed into the safety tube using a one handed technique.

- 3.9.3. Forensic Chemistry will follow the official SDPD Procedures on handling currency in impounds. The policies are outlined below, see SDPD Procedures
 3.02 Investigations and 3.15 Investigations for the most updated versions.
 - 3.9.3.1. All money is to be entered in the designated fields for currency and coins, with the total amount impounded entered in the "Money Total Value" field. Do not enter a dollar sign (\$), only the amount.

3.9.3.1.1. Cash amounts of **\$20.00 and over** will be stored separate from other impounded property.

3.9.3.1.2. Cash amounts of **less than \$20.00** should be placed in an envelope but can remain with the other impounded items.

Currency that contains possible drug residue will not be separated from the other drug items.

If narcotics are rolled into a dollar bill (any denomination) and used as a tooter, snorter, etc., or the money has any visible narcotic residue, it must be impounded in the EvidenceOnQ database. When the impound is up for disposal, any money with narcotic substance or narcotic residue will be destroyed per vault policy.

- 3.9.3.3. Documenting the serial numbers of all currency in the case notes is recommended.
- 3.9.4. Impounds annotated "Hold for grints" of "Hold for DNA" will be handled in a manner to preserve possible fingerprines and prevent DNA contamination.
 - 3.9.4.1. Criminalists will wear a freeh pair of gloves when working with these cases to prevent the possibility of depositing their prints or DNA on items.
 - 3.9.4.1.1. Cotton liners can also be worn under the gloves to further prevent the deposit of prints.
 - 3.9.4.2. Criminalist should wear masks while working cases marked "Hold for DNA."
 - 3.9.4.3. Items that may be processed for fingerprints should be handled as little as possible and in areas generally not suitable for print processing. The Criminalist should handle the evidence carefully to prevent the obliteration of possible prints.
 - 3.9.4.4. Paraphernalia such as pipes or spoons are generally not examined. They can be left in the impound envelope without processing or analysis.
 - 3.9.4.5. These cases do not need special repackaging at this stage.
- 3.9.5. When the clerical staff receives requests to process seized drug evidence for fingerprint or DNA processing, they will annotate the need for separation by

5.9.3.2.

.2.1.

the Forensic Chemistry Unit on the top of the request and forward a copy of the request to the unit supervisor.

- 3.9.5.1. The Criminalist will obtain the impounds needing separation from the Narcotics Vault.
- The Criminalist will remove and repack any suspected 3.9.5.2. controlled substances from the items of interest into appropriate containers annotating that it is a repack and labeling them with the barcode, date, and their initials. The Criminalist will repack the original packaging to be 3.9.5.3. processed for fingerprints into a second package and annotate that it is the original packaging with the barcode, date, and initials. Other items that may be suitable for fingerprint processing should also be placed in this second package. The crime scene unit can be contacted if the Criminalist has any questions about items suitable for fingerprint processing. Once the original packaging has been repacked for latent print processing, a new barcode must be generated in the videnceOnQ system. The following procedures can be used:
 - 3.9.5.1. Open the desktop version of the EvidenceOnQ program and type the incident number into the incident number field.
 3.9.5.2. Select "New" or "Clone" New will generate a new
 - 2. Select "New" or "Clone" New will generate a new barcode with the original case information automatically filled in. Clone will generate a new bar ode with all information duplicated.
 - 3.9.5.5.3. Fill in the appropriate information for the newly created item, including the item type, date and time the item was generated, and the officer that recovered the original item.
 - 3.9.5.5.4. The "Additional Description" field must be annotated with h formation indicating the originating barcole number and a brief description of the newly generated item.
 - 3.9.5.5.5. A new barcode label must be printed and attached to the outside of the new imported envelope. Use the "Print Barcode Label" by a m to print labels.
 - 3.9.5.5.6. All items will be returned to the Vaul.
- 3.9.5.6 The Criminalist will edit the annotation at the top of the request to indicate that the items have been repacked, and will include the new barcode number, date and their initials.
- 3.9.5.7 A copy of the request will remain with the case packet as at notes page. The original will be placed in the crime scene unit s mail bin for processing.
- 3.9.5.8 The Criminalist will not repack large seizures of controlled substances for processing.

4.0 POLICIES

4.1 SEIZED DRUG ANALYSIS

- 4.1.1. Only one case shall be open in the Criminalist's work area at a time.
- 4.1.2. All impounds will be thoroughly inventoried.

Each Criminalist must determine the appropriate tests to use based on the type of suspected drug.

- 4.1.4. If after performing color or instrumental tests a Criminalist decides not to conducc further testing on an item, provided that a controlled substance is being reported for that incident and individual, the Criminalist may write "initial examonly" (IEO) in their notes and stop testing. This does not need to be written if the printed report. This can also be done with federally controlled et near-controlled medication with a visual inspection or preliminary identification as well as marijuana and its products. If it is the only item, the report will reflect the apparent visual identification or test performed. No further analysis will be done.
- 4.1.5. Color tests may be done in the main hood or at the Criminalist's laminar flow station. Crystal tests may be done at the Criminalist's laminar flow station.
- 4.1.6. The base form of cocaine is distinguished from the salt.

4.1.6.1. The base form is distinguished during preliminary examination in one of the following way *

4.1.6.1.1.	An appropriate wag ner color test

- 4.1.6.1.2. FTIR
- 4.1.6.1.3. Raman
- 4.1.6.1.4. GCMS in Hexanes with one of the above

*note: the physical form (waxy/rock-like verses powder/compressed powder) of the sample while not sufficient for identification, can be useful information to direct testing and understand results.

- 4.1.6.2. A second confirmatory test must be performed in, addition to the preliminary testing above, to report the base form for the Final report.
- 4.1.6.3. If the base form cannot be confirmed the sample will be reported as cocaine, not as cocaine base due to cocaine base being a higher schedule.
- 4.1.7. Weights will be taken of all solid, non-tablet/capsule, samples analyzed and reported, with the exception of NCSDs .

- 4.1.7.1. Gross weights can be taken for any preliminary case that is not a felony drug case.
- 4.1.7.2. Gross weights can be taken for all fentanyl cases regardless of felony or court status.
- 4.1.7.3. Net weights will be taken for all felony drug cases and all court cases.

All weights taken on Criminalist's balances will be taken to two decimal places.

Individual weights of \leq 200 grams may be taken on Criminalist's balances.

All weights taken on Criminalist's balances must be taken dynamically unless the sample consistency prevents it. Note: Dynamic weighting means transferring the ontents of the item directly on to the balance and noting the weight. Static weighing refers to being able to tare a weigh boat on paper, remove it from the balance to add contents of the item to it, then returning to the balance for a weight.

- 4.1.7.7 Individual weights > 200 grams must be taken on the bulk balance.
- 4.1.8. If other items are present in a submission that could be controlled but were not examined, that should be made clear in the notes.
- 4.1.9. If an item is only being weighed for a defective or attorney, this will be documented in a printed email or using a communication log. If there is no analysis on the item, a report does not need to be created.
 - 4.1.9.1. The added weights must be technically reviewed. This can be done in the notes next to the data
 - 4.1.9.2. Communication documentation must be administratively reviewed. This can be documented next on the communication log or printed email.
- 4.1.10. Volumes of tested liquids will be estimated and documented in the notes but will not be listed on reports.
- 4.1.11. Criminalists may only issue reports in areas, and using testing methods, for which they have been approved for independent casework.

4.1.11.1.	subst	riminalist begins a case and determines that the ances present may be outside the scope for which they are oved, they must:
4.1.1	1.1.1.	Stop testing
4.1.1	1.1.2.	Document the reason for stopping in their notes
4.1.1	1.1.3.	Find a qualified criminalist to rework the case and

document the Criminalist in their notes

4.1.7.4.

- Seal and return the evidence to the Narcotics Vault. 4.1.11.1.4.
- Transfer their notes and documentation to the qualified 4.1.11.1.5. Criminalist
- The qualified Criminalist will obtain the case from the Narcotics 4.1.11.2. Vault and rework it.
 - 4.1.11.2.1. The original notes from the first Criminalist will be maintained in the new case packet.
 - If the Criminalist has already begun work on additional items in the case that are within the scope for which they are approved, they may continue testing and reporting on those items.
 - If it's possible without creating a new barcode, the items for which they are not approved to do casework should, be separated from the other items and from the original outer packaging and repackaged separately into new appropriate outer packaging.
 - .1.11.3.1.1. This separation will be documented in the Criminalist's notes.

These items will then follow steps 4.1.11.1.1.-4.1.11.1.5. The qualified Criminalist in these cases will rework only these items and will produce a Supplemental Report. The work done by the first Criminalist must be acknowledged in the subsequent Criminalist's note

ACCEPTABLE CRITERIA FOR PAPARIM NARY REPORTS 4.2

preset

- 4.2.1. Preliminary testing will be performed or poyders, liquids, solids, unidentified pills, etc. for use at preliminary hearings. If a case goes on to trial, a final report will be prepared.
- 4.2.2. In general, most cases, with the exception of tharp acoutical preparations and odd drugs, will be preliminarily analyzed by at least two-independent tests. These include a combination of two of the following

4.2.2.1. Color tests

4.1.11.3.

4.1.11.3.1.

4.1.11.

4.1.11.3.4.

- Crystal tests 4.2.2.2.
- 4.2.2.3.Instrumental tests
- Microscopic examination (mushrooms) 4.2.2.4.
- 4.2.3. Preliminary analysis can consist of only instrumental testing without microscopy, or color and crystal tests.
- The Criminalist may only report out results for those items that have been 4.2.4. analyzed.
- 4.2.5. All reports will be technically and administratively reviewed prior to release.

ACCEPTABLE CRITERIA FOR FINAL REPORTS 4.3

- 4.3.1. All final examinations, unless specifically exempted in this manual, will require a minimum of two confirmatory tests (an identifying crystal test and instrumental test or two instrumental tests).
 - If standards are not available for GC retention time comparison 4.3.1.1. without another instrumental test, and a crystal test is not available for identification, the MS data alone can be used for identification. If identification of an unknown substance is made using only the MS data, the final result will be reported as "MS only" indicating that a retention time identification was not made.
- The Criminalist may only report out results for those items that have been 4.3.2. analyzed.
- 4.3.3. All reports will be technically and administratively reviewed prior to release.

MINIMUM TESTS FOP "NO CONTROLLED SUBSTANCE DETECTED" (NCSD) 4.4 RESULTS

4.4.1. The minimum battery of tests to be performed on substances to determine that no controlled substance was refected includes, but is not limited to, the following:

4.4.1.1	Tests	s performed ar	base	d on the for	rm of the unknown.	
	4.4.1.1.1	Solids:				
		Wagner's		N		
		Marquis				
		Cobaltous Th	niocva	nate		

Nitroprusside Gold Chloride (Crystal Test Stereoscopic exam NJ CSA Duquenois-Levine (if dark subs Ferric chloride

4.4.1.1.2 Powders: Same as above, plus: Mecke

Liquids: 4.4.1.1.3 Wagners Marquis Cobaltous Thiocyanate Nitroprusside Libermanns Duquenois plus Chens 2 Ferric Chloride

4.4.1.1.4 Other:

- 4.4.1.1.4.1. If the material appears to be something such as soap or nuts, the Criminalist will describe the material in their notes and report as "apparent ...," and proceed with chemical testing if necessary. If chemical testing leads the Criminalist to conclude that no controlled substances are present, then "No controlled substance detected" will be reported.
- 4.4.1.1.4.2. If the material cannot be cut with a knife or does not appear suitable for chemical testing, such as a stone or piece of glass, the Criminalist will describe the item in their notes and report as "apparent ..." and:

If **no testing** was attempted, "Item ... is not suitable for analysis and was not laboratory examined," will be reported.

If **microscopy** was conducted, the test performed will be included in the report. "Item ... was determined by microscopy not to be suitable for analysis. No further analysis was conducted," will be reported.

4.4.1.2. A colored liquid may interfere with the color tests for GHB/GBL/1,4,4 Burnediol. Those samples will be analyzed either by a combination of GCMS and FTIR, or by GCMS alone with the inclusion of a standard.

- 4.4.2. Color and/or crystal tests do not need to be performed to conclude that no controlled substances were detected if the sample is analyzed by GCMS using the universal program. Analysis by FTIR of Ray and an be used to conclude that no controlled substances were detected if analysis by FTIR or Raman identifies the presence of a non-controlled substance
- 4.4.3. Mushroom material must be soaked overnight and run or GCMS prior to an NCSD determination. See Botanicals section.
- 4.4.4. Samples containing acetaminophen must be extracted or run on prappropriate GCMS method to remove acetaminophen from the analytical results to ensure no controlled substances are present prior to an NCSP determination.
- 4.4.5. If the results of testing are negative, or do not indicate the presence of a controlled substance, the report will read, "no controlled substance detected

4.5 CONSUMING SAMPLES FOR ANALYSIS

4.5.1. Occasionally, consuming a sample during analysis is required. In these instances, the unit supervisor is notified. In addition, permission to consume the sample should be obtained from the attorney assigned to the case or, if an

Nor

attorney has not been assigned to the case, the detective assigned to the case. Three business days will be allowed after the Criminalist has reached out to the attorney or detective before proceeding with evidence consumption in the absence of a response. This process should be documented in the case notes.

4.5.2. After consuming the samples, Criminalists should save any remaining extracts. These extracts would be available should additional work be required by the original Criminalist, another Criminalist if necessary, or a defense expert. Extracts can be placed in crimped GC/MS vials and maintained with the original impound. Notes will indicate how the sample was prepared and maintained. At a minimum, all extracts will be labeled with the incident number, item number, and initial, or the barcode number and initials.

4.6 MARKING ANALYZED ITEMS

- 4.6.1. Indicidual containers housing analyzed items must be identified to include:
 - 4.6.1.1Barcode number4.6.1.2Lutruls
- 4.6.2. If multiple solid substances are housed in the same container, the Criminalist must repack or put some marking on the specific substance analyzed.

4.7 REQUESTS FOR EVIDENC

- 4.7.1. The laboratory will comply with court orders for release or splits of evidence.
- 4.7.2. Samples will not be released until a mail aboratory analysis has been completed, unless it is to another law enforcement agency.
- 4.7.3. Whenever possible, the original Criminalist whenever possible for release.
- 4.7.4. When splitting a sample for release, the Criminalist must tenerate a new barcode in the EvidenceOnQ system for the newly generated simple following the procedure outlined in section 3.9 of this manual.
- 4.7.5. The case packet will be annotated indicating the weight of the meterial prepared, the incident number, barcode, date and initials of the Criminalist. A copy of the court order will be attached to the case packet.
- 4.7.6. The item to be released, and the copy of the court order received, will be turned in to the Vault for release.

4.8 ACID NEUTRALIZATION PROCEDURE

4.8.1. During analysis of seized drugs, small amounts of acid are generated in spot wells on plates. The plates are placed in a stoppered sink with water and sodium bicarbonate. Disposable plates may be placed in a neutralizing jar in a laminar hood prior to being placed in the sink or disposed of.

- 4.8.2. Prior to discharge into the sewage system, the water solution will be checked with pH paper to ensure a neutral pH (pH range of 6.0–9.5). The minimum safety equipment worn by the Criminalist or technician neutralizing the acid or washing spot plates includes gloves, safety glasses, and a lab coat.
- 4.8.3. A neutralization log will be kept to record the date, operator initials, type of waste treated, approximate amount, and the pH determined after treatment.

.9 CRYSTAL TEST WASTE

4.9.1. Was e generated by crystal testing will be placed in a sharps container that has had the biolog and labels defaced. These containers will be taken to the Narcotic Vaul, when full.

4.10 FENTANYL CLEAN-UP

- 4.10.1 Wear appropriate personal protective equipment (PPE).
- 4.10.2 Add 1 teaspoon full of power exiclean to 500 mL of water in a spray bottle. Shake gently until all powder is in solution.
 - 4.10.2.1. Completely cover any potentially contaminated area with spray.
 - 4.10.2.2. Within 15 minutes, scrub well a paper towel until dry. Do not let the solution evaporate.
 - 4.10.2.3. Dispose of paper towels and disposable PPE in a biohazard bin.

4.11 GAS SUPPLIES AND ROOM 138 SUPPLIES

- 4.11.1. The laboratory technician will ensure Room 138 is stocked for use. Supply requests, including lab-wide supplies, will be processed through the laboratory technician responsible for ordering supplies.
- 4.11.2. The gas delivery truck driver brings filled compressed gas tanks to the Police Department and removes the empty tanks. The tanks are currently stored if the Sally Port on the first floor. The laboratory employee that meets the criver and escorts him/her into the building will be responsible for signing the invoice and providing a copy of the invoice to the clerical staff.

5.0 CASE DOCUMENTATION

5.1 NOTES

- 5.1.1. The note pages will be numbered. The total number of pages will be annotated on the first and last pages of the notes. If subsequent pages are added to a packet, total number of pages will be updated accordingly.
 - 2. All note pages will contain the incident number, Criminalist's initials, page number, and the date. Barcodes are used to identify items within the note pages. Abbreviated item identifiers may be used instead of barcodes, if a key clearly associates those numbers with the barcodes.

5.1.2.1. For case types other than Seized drugs (ex: homicides), the case humber must be included on each note page.

- 5.1.3. The report and case notes will be maintained with the laboratory case files.
- 5.1.4. Notes must be legible abbreviations must be common and understandable, or listed in the approved abbreviation list (see section 17); permanent ink must be used.
- 5.1.5. Corrections and inserted notes will be initialed. If the corrections or interlineations are done on a loce ones than the date listed on the report or on the note page, the correction/interlineations(s) will be dated.
- 5.1.6. Reference materials and sources relied on to form conclusions will be noted and included. This includes the instrument library spectra, Drug I.D. Bible edition and page, pill identifier, etc.
- 5.1.7. Evidence disposition must be listed in notes.
- 5.1.8. Notes must include the start and end dates of analysis
- 5.1.9. All barcodes received will be listed in the notes. Descriptions or all items analyzed will be included.
- 5.1.10. Any weights, measurements, or estimated volumes made must be included in the notes.
 - 5.1.10.1. Material may be described as "apparent trace" in notes based in the amount of material present.
 - 5.1.10.1.1. Any material weighing less than or equal to the reported uncertainty of measurement is reported as trace.
 - 5.1.10.2. Residue is a descriptive term, not a qualitative one, and will not be used in place of a recorded weight.
 - 5.1.10.2.1. Residue refers to material in an item that is not conducive to collection for weighing and testing.

Examples: melted material in a pipe, traces of powder that cannot be removed from a bag, material thinly smeared on foil.

- 5.1.11. All testing conducted must be listed in the notes and all test results used to form conclusions must be included.
- 5.1.12. Conclusions will be included in the notes.
 - 1.13. Documents such as work requests and communication logs will be included in the case packet as notes. All pertinent case information and date of inclusion must be present.

1.14. Communications affecting testing, or giving opinions or results beyond those already released, must be documented.

- 5.1.1.1.1 The Criminalist may document the communication via a printed email added to the case packet, by writing it into the case notes bages, or by using a communication log.
- 5.1.14.2. The occumentation must include who the communication was between, the date, a brief description of the topics or results discussed, and any decisions made during the communication.

5.2 REPORT FORMAT

- 5.2.1. Criminalist's conclusions are typically intered into the Narcotics Database prior to generating a written report. A Dicrosoft Word report template, is then used to generate the report.
 - 5.2.1.1. Reports can be hand typed using template when necessary (ex. Database is offline, reports for internal Affairs).
- 5.2.2. Each report must include:
 - 5.2.2.1. The last name/identifier of the defendant() listed on the barcodes analyzed
 - 5.2.2.2. Incident number
 - 5.2.2.3. Incident type
 - 5.2.2.4. Criminalist's name and PD ID number
 - 5.2.2.5. Arrest/incident date
 - 5.2.2.6. Requesting Officer
 - 5.2.2.7. Barcode numbers of the items analyzed
 - 5.2.2.8. Packaging information, including the type of containers

- Descriptions of the items reported 5.2.2.9.
- Weights (to two decimal places) or measurements, if applicable, 5.2.2.10. and the associated U of M at k=3.
 - If a total weight of items is being reported, the U of M 5.2.2.10.1. statement must indicate that the U of M is per each weight taken.

Static weights taken on Criminalist's balances must 5.2.2.10.2. include the corrected U of M (twice the U of M of a dynamic weighing) and must be documented in the notes.

- 5.2.2.10.3. Since samples weighing less than 0.04 grams are not routinely tested, these items only need to be listed on the report if nothing else was tested for that suspect.
 - 5.2.2.10.3.1. Weights of less than or equal to the reported uncertainty of measurement will be reported as "trace" when included on reports.

Volumes are not reported

kan inations performed on the reported items 5.2.2.11.

- Opinions and interpretation 5.2.2.12.
- Date of authorization 5.2.2.13.
- Initials and cate of technical and administrative reviewers and 5.2.2.14. date of issuance
- 5.2.3. If multiple items are examined, the report must specify which items are included in the color, crystal, visual, and instrumental testing.
- 5.2.4. When weights are reported, the footer of herefort and will indicate the confidence level used.
- 5.2.5. Disposition of evidence will be included on the report
- 5.2.6. Example report header:

SUSPECT: INCIDENT #: INCIDENT TYPE: INCIDENT DATE: OFFICER: CRIMINALIST:

DOE, JOHN XXXXXXXXXXXX NARCOTICS

- L'USA 5.2.7. Reports for investigative units other than Narcotics will need the following additional information added to the header:
 - Victim 5.2.7.1.
 - 5.2.7.2. Case number

5.2.7.3. Detective, if listed, in place of the Officer

5.2.7.4. Charge

- 5.2.8. The title of the report will state if the report is Preliminary, Final, or Supplemental.
 - 5.2.8.1. Supplemental worked done to produce the first Final report will not be titled as supplemental.
 - 5.2.8.2. Supplemental work done to produce additional Preliminary or additional Final reports will be titled as Supplemental reports.

DISTRIBUTION AND RETENTION

- 5.3.1. Reports are faxed or emailed on a regular basis to their appropriate end users by the derical staff.
- 5.3.2. Final reports are faxed or emailed following administrative review to the District Accorner's Office or City Attorney's Office.
- 5.3.3. Original report packets will be filed in the Narcotics Files located in the laboratory. The clerical staff, following faxing or emailing of the reports, will file case packets.
- 5.3.4. Case packets are scanned to be riled electronically on the LAN. Hard copies are filed after scanning and will be maintained for the current year plus the previous 2 years; only the electronic file will be kept after this time period.
- 5.3.5. Requests for copies of reports will be referred to the clerical staff.
- 5.3.6. Defense attorneys will be referred to the prosecutor's office for copies of reports involving criminal cases.
- 5.3.7. Requests for copies of reports for civil cases will be referred to the unit supervisor.
- 5.3.8. Requests for reports by other agencies will be referred to the narcotics detective to avoid possible conflict with criminal investigations.

5.4 NARCOTICS DATABASE

- 5.4.1. Each impound must be imported into the Narcotics Database from the EvidenceOnQ database by the case criminalist. Impounds are imported using the following steps:
 - 5.4.1.1. Open the Narcotics Database.
 - 5.4.1.2. Click the "Scan Barcode" button.
 - 5.4.1.3. Type or scan the barcode numbers of the items to be reported, this can be done individually or in a batch.

5.4.1.4. Select "Import." Close the window by selecting "Return" after the hourglass disappears.

5.4.2. To enter data into the database

- 5.4.2.1. Type or scan the barcode of one of the items. All previously entered items for that incident number should populate.
 - 5.4.2.2. Click on the barcode item line and then click edit. 5.4.2.2.1. If the item was previously reported, click the barcode item line and then click "Court analysis" and continue as listed below.
 - Add all necessary information and click "save and return," Or, if more items need to be entered under the same barcode, click "save and add next," enter the information, repeat as necessary, and then click "save and return."

5.4.2.4. Repert for all items to be reported.

- 5.4.3. Generate report
 - 5.4.3.1. Select alking lines to be reported and click "Forensic Report."
 - 5.4.3.2. Edit generated report as needed.
 5.4.3.2.1. Verify an necessary information is present and correct in the header.
 5.4.3.2.2 Ensure that deprivations are correct and complete.
 - 5.4.3.2.2. Ensure that accorrigations are correct and complete.
- 5.4.4. Reviewing and releasing results
 - 5.4.4.1. After conducting a technical review or the case packet, the reviewer will review and release the data entered into the Narcotics Database.
 - 5.4.4.1.1. Click "Bar Code" or "Incident Number" and type in the appropriate number.
 - 5.4.4.1.2. Ensure that all provided header information matches the report and notes.
 - 5.4.4.1.3. For each item click the line of that item then click "Edit" and ensure that all of the item testing information matches the report and notes.
 - 5.4.4.1.3.1. Click "Return" after reviewing each one.
 - 5.4.4.1.4. For each item, if the information is correct, click the line of that item and then click "Tech Review."
 - 5.4.4.1.5. For each item, if the information is correct, click the line of that item and then click "Admin Review." This will release the results.
 - 5.4.4.1.6. If any information is incorrect, work with the Criminalist who did the work to correct the issues before clicking on "Tech Review."

5.5 STATISTICS

5.5.1.2.

5.5

- 5.5.1. Each Criminalist reports their daily activities on individual monthly stat sheets due to the unit supervisor by the 5th workday of the following month. Seized drug stats will include:
 - 5.5.1.1. Number of preliminary cases examined, and the number of samples analyzed
 - Number of court cases examined, and the number of samples analyzed
 - Number of court appearances and hours of court time, including court preparation
 - Number of HQ cases not analyzed within 24 hours
 - 5.5.1.5. **Fraining time**
 - 5.5.1.6. Special projects
 - 5.5.1.7. Review nours
 - 5.5.1.8. Narcan continue
 - 5.5.1.9. Meetings
 - 5.5.1.10. Crime Scenes and Grime scene reports
 - 5.5.1.11. Other activities should also be reported
- 5.5.2. For clarification:
 - 5.5.2.1. If six plastic bags are described buy only three are analyzed, this constitutes three items for statistical purposes.
 - 5.5.2.2. If color, crystal, and GCMS were conducted on all three items, this is still three items tested.
 - 5.5.2.3. One written report represents one case.

C Ser

6.0 EQUIPMENT

6.1 SEIZED DRUG EQUIPMENT LIST

- 6.1.1. The Forensic Chemistry Unit utilizes the following items of equipment (see chart below for specifics):
- <u>GCMS:</u> for rapid separation of compounds. 6.1.1.1. <u>Polarized Light Microscope</u>: For monitoring the various stages 1.1.2. of crystal growth during a microcrystalline test and observing spores. Minimum magnification required: 100x. <u>Stereomicroscope:</u> The stereomicroscope is used for examining 6.1.1 plant structures. Low power magnification needed (4-40x pproximately). En st onic Balance: For the weight determination of apparent 6.1.1.4. drug substances and in making reagents. Incubition Oven: For drying or catalyzing chemical reactions 6.1.1.5. through the addition of heat. <u>Fume Hood:</u> Hrowides a safer environment by providing a place to work with characteristic color tests, extractions, etc.). 6.1.1.6. FTIR: For rapid identification of drug compounds and their 6.1.1.7. isomers. 6.1.1.8. Raman: For rapid identification of drugs, including some mixtures, and ones contained in packaging. Refrigerator/Freezer: To store standar is, samples, and 6.1.1.9. solutions. 6.1.1.10. Hot plates: For heating and/or stirring solutions. UV box: For testing substances for UV reactivity. 6.1.1.11. C SA 6.1.1.12. Vortex: To rapidly mix solutions.

Equipment	Make	Model	Serial#	Software version
	Aligent			MSDChemStation
GCMS-75 (GC)	Technologies	G3442B	CN1433073	E.02.00.493
	Aligent			or higher
GCMS-75 (MS)	Technologies	G7038A	US1424L215	
	Aligent			MassHunter GC/MS
GCMS-77 (GC)	Technologies	G3440A	CN10826025	Acquisition B.07.01.1805
GCMS-77 (MS)	Aligent	Carra	11004020550	or higher
GCMS-77 (MS)	Technologies	G3171A	US81829559	0
FTIP	Thermo Scientific	Nicolet iS10	AKX1200568	OMNIC 8.3.103 or higher
			AKA1200508	
Raman	Thermo Scientific	DXR Raman	AI21200569	OMNIC 8.3.104 or higher
Microscopes	Nikon	Labphot	955129	_
Microscopes 2	Leira	DM750P	D540313366CW0018	_
		Optiphot-		
Microscopes 3	NIK N	POL	142522	-
Microscopes 4	Leica		D540309307CV0034	-
DG :	Nilson	Optiphot-		
Microscopes 5	Nikon	POL Letipse	141301	-
Microscopes 6	Nikon	1400 DOL	531346	-
Microscopes 7	Leica	D'17 '0'	D540307732CU0108	
Stereoscope 1	Olympus	SZ3	SZ3060	_
Stereoscope 2	Leica	S6E	1446339	_
Stereoscope 3	Leica	MZ95	143)09	_
Balance 1	Denver	S-403	5050/44	-
Balance 2	Denver	S-403	25057037	_
Balance 3	Mettler	PE24	E882/0	-
Balance 4	Denver	S-403	25050643	-
Balance 5	Denver	S-403	25050640	-
Balance 6	-	_		-
Balance 7	Denver	S-403	26450329	-
Fridges	Thermo			
(Narc/Reagent)	Scientific Fisher	10ECEETSA	0158391701160331	-
hot/stir plates	Scientific	Isotemp	C3720421091648703	
, 1	Fisher	· · ·		
hot/stir plates	Scientific	Isotemp	C3720421091648628	
oven (in breath	Thermo			
room)	Scientific	PR305220M	300100699	_ •
Ductless Hood	Air Science	PureAir		-
Long Wave UV 365nm Box	Spectroline	ENE-260C	2073607	

	Scientific Industries,			
Vortex	Inc.	G-560	2430809	

6.2 GCMS PERFORMANCE CHECKS

6.2.1.1.

5.2.1.2

6.2.1. A STANDARD or AUTOTUNE is performed, and evaluated, weekly when the instrument is in use and after any maintenance that directly effects separation or identification is performed.

- QUICKTUNES and TARGET TUNES can be performed at the Criminalist's discretion.
 - Peak widths should be between .45 to .65 amu.
- 6.2.3. Peaks should be smooth and symmetrical.
- 6.2.1.4. Mass peaks should be +/- 0.2 amu
- 6.2.1.5. ENVolts approaching 3000 may indicate that source cleaning is needed.
- 6.2.1.6. N_2 and H_2O relative abundances should each be less than 10%.
- 6.2.1.7. Relative abundances should be within the following limits: 69.00 70 - 100% 219.00 >30%* 502.00 > 1/0 * 210.00 may be greater them 0.00% if it's permalized to the

* 219.00 may be greater t'an 100% if it's normalized to the 69.00 peak.

6.2.1.8. Iso Ratio: Isotope Mass Ratios should be within the following limits: 70.0 0.5 - 1.6%

70.0	0.5 - 1.6% 🤞
220.0	3.2 - 5.4%
503.0	7.9 - 12.3%

- 6.2.1.9. Tune reports are kept on file, in chronological order, in binders kept near the instruments.
- 6.2.1.10. The Criminalist or laboratory technician who evaluates me tune will initial it.
- 6.2.1.11. If the result of the tune does not meet acceptable criteria mentioned above, no casework will be conducted until the tune problem is resolved. See section 7.2.
- 6.2.2. Quarterly, a mix of laboratory standards with close elution times along with low and high molecular weights will be run on the universal method.
 - 6.2.2.1. This system check will ensure that the retention times of related compounds can be separated and each component identified.

- 6.2.2.2. The results will be evaluated as per 6.2.2.1 by a Criminalist and initialed before being filed in the instrument binder.
- 6.2.2.3. If the result of the check does not meet acceptable criteria, no casework will be conducted using that instrument until the problem is resolved. See section 7.2.
- 6.2.3. The GCMSs are covered by outside vendor service contracts for repair and maintenance.

. Unit Criminalists and laboratory technicians can conduct periodic cleaning and maintenance of the GCMSs when needed.

6.2.5. An outside vendor will perform preventative maintenance of the instruments on an annual basis.

6.2.5.1. Labels affixed to the instrument will indicate the date of last preventative maintenance and due date of the next.

6.2.6. Problems, manupunce, etc., are documented in the individual instrument maintenance binder located in the GCMS room.

6.3 FTIR CALIBRATION AND PLRFORMANCE CHECKS

- 6.3.1. Quarterly, and after any maintenance is performed, a VAL-Q calibration check will be run.
 - 6.3.1.1. The check is automated, gastre that all criteria read "PASS"
 - 6.3.1.2. A Thermo Fisher polystyre e standard will then be run under the same conditions as evidentity samples.

6.3.1.2.1. The polystyrene standard will be treated as an unknown and evaluated and compared to the libraries as per sections 12.2 and 12.3 of this manual.

- 6.3.1.3. The results will be evaluated as per sections 11.2 and 12.3 by a Criminalist and initialed before being filed in the instrument binder.
- 6.3.1.4. If the result of the check does not meet acceptable criteria, acceptable criteria, acceptable criteria, acceptable criteria, acceptable will be conducted using that instrument until the problem is resolved. See section 7.2.
- 6.3.2. The FTIR is covered by an outside vendor service contract for repair and maintenance.
- 6.3.3. An outside vendor will perform preventative maintenance of the instrument on an annual basis.

- 6.3.3.1. A label affixed to the instrument will indicate the date of last preventative maintenance and due date of the next.
- 6.3.4. Problems, maintenance, etc., are documented in the individual instrument maintenance binder located near the instrument.

6.4 RAMAN CALIBRATION AND PERFORMANCE CHECKS

 Alignment and calibration of the laser on the Raman must be performed every 30 days or less, and after any maintenance is performed, using the calibration platform.

The checks are automated and a Criminalist or laboratory technician must ensure that each has passed.

- 2. One of the 30 day alignments and calibrations should be additionally recorded as a quarterly check.
- 6.4.2. In addition to the 30 day alignment and calibration, a Thermo Fisher polystyrene standard will be run quarterly, and after any maintenance is performed, under the same conditions as evidential samples.
 - 6.4.2.1. The polystyrere standard will be treated as an unknown and evaluated and compared to the libraries as per sections 13.2 and 13.3 of this manual.
- 6.4.3. The results for the quarterly checks will be evaluated by a Criminalist and initialed before being filed in the instrument binder.
- 6.4.4. If the results of the checks do not meet acceptable criteria, no casework will be conducted using that instrument until the prebrum is resolved. See section 7.2.
- 6.4.5. The Raman is covered by an outside vendor service contract for repair and maintenance.
- 6.4.6. An outside vendor will perform preventative maintenance of the instruments on an annual basis.
 - 6.4.6.1. A label affixed to the instrument will indicate the fate of last preventative maintenance and due date of the next.
- 6.4.7. Problems, maintenance, etc., are documented in the individual instrumen maintenance binder located near the instrument.

6.5 BALANCE AND WEIGHT CALIBRATION CHECKS

- 6.5.1. Calibration checks will be performed on a quarterly basis, and after any maintenance is performed using NIST traceable standard weights.
 - 6.5.1.1. Checks will be performed to three decimal places.

- 6.5.1.2. Worksheets will be completed by the laboratory technician or Criminalist performing the checks.
- 6.5.1.3. Weights must fall within the current specified Uncertainty of Measurement for the balance being checked to be acceptable.

6.5.1.4. Following review by the unit supervisor, completed worksheets will be maintained in the Maintenance binder, which is kept in the Forensic Chemistry Unit.

6.5.1.5. If the results of the checks do not meet acceptable criteria, no casework will be conducted using that balance until the problem is resolved. See section 7.2.

6.5.2. An outside vendor will perform calibration and maintenance of the balances on an annual basis.

6.5.2.1. A label affixed to the balance will indicate the date of last alibration and due date of the next.

- 6.5.2.2. **The Quality Assurance Manager will make arrangements for the** outside service.
- 6.5.3. The NIST traceable standard weights will be calibrated every four years by an outside vendor.
 - 6.5.3.1. The outside vender must be accredited by an accrediting body subject to ILAC.
 - 6.5.3.2. The quality assurance manager will make arrangements for the outside service.
 - 6.5.3.3. The weights will then be checked annually after balance calibration.
- 6.5.4. Uncertainty of Measurement evaluation will be done if there is any maintenance done on a balance that would affect weighing capability, a new balance is purchased, or a new Criminalist starts in the Unit
 - 6.5.4.1. The maximum calculated repeatability and linearity measurements along with manufacturer specifications and vendor calibration data are evaluated for incorporation into the calculation of combined standard uncertainty.
 - 6.5.4.2. Expanded uncertainties are calculated for both the 95.45% and 99.7% confidence intervals.
 - 6.5.4.3. Measurements will be taken using NIST-traceable weights.

6.5.4.3.1. For analyst's benchtop analytical balances, a minimum of ten replicates of each 0.010 g, 1.000 g, 10.000 g, 50.000 g, and 100.000 g; and 20 replicates of each 0.020 g, 2.000 g, and 20.000 g will be taken.

- 6.5.4.3.2. For the bulk balance, a minimum of three replicates, on different days at different times of day, of each of the five larger weights (100g-1Kg) will be taken.
- 6.5.4.4. The maximum standard deviation from all of the weights on all of the analytical balances is used in the uncertainty of measurement calculations.
- 6.5.4.5. Additionally, for analyst's analytical balances, over the course of five days (not necessarily consecutive), criminalists perform measurements using a set of NIST traceable weights on their balances with four different target measurements.

6.5.4.5.1. The four measurements are 0.050 grams, 5.000 grams, 40.000 grams, and 200.000 grams.

Criminalists will do individual placement of the weights needed to achieve the target measurement. Criminalists will also place the weights in different

positions on the weighing pan.

Measurements will be taken in both the morning and the afternoon.

6.5.4.6. The calculated standard uncertainties are plugged into the following formula:

$$U_c = \sqrt{u(x)^2 + u(y)^2 + u(z)}...$$

6.5.4.5.2.

- $U = k \ast_{u_c}$ Where U is the expanded uncertainty and k is the coverage factor
- 6.5.4.7. All measurements will be reptin the uncertainty binder, in the lab, including the worksheets generated to record balance measurements.
- 6.5.4.8. When a new criminalist is added to me unit, the uncertainty of measurement will be determined on their analytical balance to confirm that the newly calculated U of M is not higher than the one currently reported. If it is higher, the reported uncertainty of measurement will be recalculated.

6.6 OTHER EQUIPMENT PERFORMANCE EVALUATION

- 6.6.1. The polarized light microscopes and stereomicroscope are all service annually by an outsider vendor coordinated by the QA Manager.
- 6.6.2. Refrigerators and freezers have NIST traceable thermometers and are checked weekly to ensure they are within established ranges.
 - 6.6.2.1. Current records are kept on the individual refrigerator and archived records will be kept in binders in the Unit.
 - 6.6.2.2. If temperatures are found to be out of range, temperature sensitive materials will be moved to another suitable location.

- 6.6.3. The UV light box is checked at time of use with UV sensitive materials.
 - 6.6.3.1. If the UV box is not working properly (ie the UV sensitive material does not fluoresce), no casework will be conducted using it until the problem is resolved. See section 7.2.
 - 6.6.3.2. Current records are kept near the UV box and archived records will be kept in a binder in the Unit.

4. Hoods are checked on a monthly basis by a Lab Safety representative.

Hoods are checked annually by an outside vendor.

6.6.5. Other equipment is repaired or replaced as needed.

6.7 USE OF LOUPMENT

66.4.1.

6.7.1. Use and maintenance of equipment will be restricted to those properly trained to do so.

6.8 REAGENT PREPARATION/TESTING

- 6.8.1. A reagent log will be maintained on all reagents used within the unit and will include:
 - 6.8.1.1. Name of the reagent
 - 6.8.1.2. Type of test it is used for
 - 6.8.1.3. Specific directions for preparation (see section 16)
 - 6.8.1.4. The test used to verify the leagent and the
 - expected results
 - 6.8.1.5. Verification test results
- 6.8.2. Each reagent will be tested by a criminalist prior to use in casework, or being placed in a criminalist's hood, with a verified standard
 - 6.8.2.1. The lot number for the new reagent will not be assigned until the reagent has been verified.
 - 6.8.2.1.1. The test date will indicate the first date of use and will be used as the lot number for the reagent.
 - 6.8.2.2. Results of testing will be recorded in the reagent log along w the initials of the Criminalist performing the test.
 - 6.8.2.3. If the expected results are not obtained during reagent verification, the reagent will not be put in to use. See section 7.2.
- 6.8.3. All reagents located in the main hood or at the Criminalist's benches will be tested on a quarterly basis.

- 6.8.3.1. Criminalists will perform the tests on color and crystal test reagents in bench hoods as well as on the color test reagents in the main hood.
- 6.8.3.2. Test results will be documented in the Color/Crystal Reagent Working Solution QC logs, which are maintained in the forensic chemistry unit reagent log binder.
- 6.8.3.3. If the results of the checks do not meet acceptable criteria, no casework will be conducted using that reagent until the problem is resolved. See section 7.2.
- .8.4. Stock bottles containing reagents will be labeled with the name of the reagent, a boundary of the reagent.
 - 6.8.4.1 Working solutions obtained from the stock bottles will be labeled with the same lot number, as well as with the name of the solution and specific hazards.
- 6.8.5. Chemical and standard containers will be labeled with the date received, and initials of the person checking them in.
- 6.8.6. Reagents housed in the main fume hood are monitored for label condition.

6.9 STANDARD PREPARATIO

- 6.9.1. A standard log will be maintained on all standards used within the unit and will include:
 - 6.9.1.1. Name of the standard
 - 6.9.1.2. Storage location
 - 6.9.1.3. Manufacturer lot number
 - 6.9.1.4. Expiration dates, if known
 - 6.9.1.5. Lab standard number
- 6.9.2. Verification of standards will be done prior to casework via intrumental analysis and manufacturer certificates, when possible.
 - 6.9.2.1. When not possible, either of the two is sufficient.
 - 6.9.2.2. The instrumental data will be evaluated as outlined in either sections 11.5-11.7, 12.2-12.3, or 13.2-13.3.
 - 6.9.2.3. Secondary drug standards may be used if they have met verification requirements
 - 6.9.2.4. Standards that do not pass verification will not be used.
- 6.9.3. Verification information and manufacturer certificates of analysis will be kept in binders labeled "Standard Verifications," located in the Forensic Chemistry Lab.

6.9.4. Verified standards will be identified by a green sticker and the letter "V" for verified. Non-verified standards will be stored in a different location.

6.9.5. Standards will be stored according to manufacturer specifications.

6.9.5.1. Standards stored refrigerated or frozen will be monitored weekly using NIST traceable thermometers.

6.9.5.1.1. If temperatures fall out of range, the standard will be verified before use. If the standard cannot be verified it will be discarded.

5.6. Standards must be labeled with the name of the standard, lab standard number, the date received or date inspected, and initials.

This does not apply to GCMS vials which must be labeled with the lab standard number at a minimum.

PLADORA, OR, USK

o.9.6.1.
7.0 QUALITY ASSURANCE

7.1 GENERAL QUALITY ASSURANCE

1.1. General Quality Assurance Policies are covered by the Quality Manual.

RERFORMANCE CHECKS

- 7.2.1. If the result of any performance check does not meet acceptable criteria, no casew rewill be conducted using that piece of equipment, reagent, or standard until the performance problem is resolved.
 - 7.2.1.1. A Quality Incident Summary Form will be filled out, if applicable (see section 7.6).
- 7.2.2. Whenever possible, the Criminalist or technician discovering the problem should attempt to treableshoot the issue while communication with the rest of the unit that the piece of ecupment, reagent, or standard should temporarily not be used in casework. This communication must be done through the use of a filled out "Troubleshooting" (ag at a minimum.
 - 7.2.2.1. If this issue is temporary for example the GCMS needs to be baked out due to high air and water after changing the liner, it does not need to be recorded in the maintenance log.
- 7.2.3. If troubleshooting fails, or the issue is persistent, the Technical Lead or Supervisor will be notified to determine if the elece of equipment, reagent, or standard needs to be pulled from service.
 - 7.2.3.1. If the equipment, solution, or standard needs to be pulled from service, this must be communicated to the rest of the unit through the use of a filled out "Out of Service" tag, at a minimum.
- 7.2.4. If the issue has potentially affected released casework results the Technical Lead and Supervisor should be notified immediately to evaluate.
- 7.2.5. All equipment maintenance, and any time a piece of equipment is removed from or returned to service, must be documented in the applicable maintenance log.

7.3 TECHNICAL AND ADMINISTRATIVE REVIEWS

7.3.1. Reports will be technically and administratively reviewed prior to dissemination following established review criteria.

- 7.3.1.1. Results can be released following a technical review.
- 7.3.2. Technical reviewers must have a current satisfactory proficiency test or be signed off in the drug category in forensic chemistry.
- 7.3.3. The reviewers will look at all technical worksheets, datasheets, and printouts within the case packet.
- 7.3.4. At the completion of their review, the reviewer will sign and date the report and the first page of the Criminalist's notes.

Administrative reviews are generally performed by the unit supervisor.

- 7.3.7. Narcotics database entries are checked and released by the technical reviewer (see section 5...4.).
- 7.3.8. The Criminalist has performed the work must address (correct or otherwise resolve) all concerns raised by the technical reviewer.
 - 7.3.8.1. Cases may not be transferred to another technical reviewer because of disagreements in the review process.
 - 7.3.8.2. If no agreement can be reached, the Criminalist will consult with the Technical Lead, together with the technical reviewer, to resolve the disagreement.
 - 7.3.8.3. Quality Incident Summary forms must be filled out, if applicable (see section 7.6)

7.4 CASE REVIEW CRITERIA

TECHNICIAL REVIEW

Performed by qualified Criminalist on all preliminary and fine reports. Name(s), incident number and barcodes (or defined identifiers) are properly

Name(s), incident number and barcodes (or defined identifiers) are properly recorded on notes and reports

Barcodes (or defined abbreviations) are used to identify items within the note pages

Evidence packaging and seals are described

Proper laboratory approved procedures were used Tests conducted or attempted and results obtained were documented

Appropriate controls, standards, and blanks were used

Supporting data, records, photos, printouts, diagrams, etc. are included

Instrument operating parameters are recorded

Criminalist's results or conclusions are reasonable, appropriate, and supported by the data, notes, and comments

Addresses all technical concerns with the Criminalist who performed the analysis. Consults with the Technical lead, together with the Criminalist who performed the analysis, to resolve any conflicts that arise during technical review as necessary

ADMINISTRATIVE REVIEW

Performed by unit supervisor or designee.

Reports are complete

All pages are numbered appropriately

Writing is legible

Notes and records are permanent (i.e. ink)

Corrections are made by an initialed single strikeout, and date if needed; no info is obliterated or erased

incident number, Criminalist's initials, and dates are on each page

r technical review has been performed by a qualified Criminalist

7.5 PROFICIENCY TESTING PROGRAM

- 7.5.1. Each criminalist signed off to do seized drug analysis must satisfactorily complete one proficiency test in seized drug analysis per calendar year.
 - 7.5.1.1. Criminalists still in seized drug training may do an increase aboratory comparison in lieu of a proficiency test.
- 7.5.2. Analysis of the samples will follow the procedures and policies used to test unknown case samples.
- 7.5.3. All samples will be taken through final analysis.
- 7.5.4. All results of proficiency and increasion tory testing must be consistent with the test provider's results to be diemed satisfactory.
 - 7.5.4.1. If the test results are unsate factory, the Technical Lead and Supervisor will assess the situation and determine the best course of action.

7.5.4.1.1. Actions may include, but are poolimited to, change in procedure, reanalysis of samples, petraining, and removal from casework.

7.5.5. Criminalists will be notified of proficiency test results via a Proficiency Test Record form.

7.6 QUALITY INCIDENT SUMMARY FORM

- 7.6.1. For any equipment failure, unexpected control result, or when a technical policy was violated in the process of analysis, a Quality Incident Summary Form (QIS) must be filled out.
- 7.6.2. QISs will be filled out by the Criminalist or Technician who discovered the issue when the issue is regarding an equipment failure or unexpected control result. When the issue is regarding a failure to follow a technical policy, the Criminalist conducting the analysis will fill out the form.

- 7.6.3. After filling out all pertinent information on the QIS, the form, along with all supporting documentation, will be submitted to the Technical Lead for tracking and any necessary follow up.
- 7.6.4. QISs will be tracked and monitored by the Technical Lead to check for trends that could indicate issues such as problems with lab equipment, training inadequacies, or process failures.
- The Technical Lead will follow up on each issue, and as 7.6.4.1. appropriate: 7.6.4.1.1. Take action to control and correct the issue. Address the consequences, to include evaluating 7.6.4.1.2. potentially effected casework. Ensure follow up action is completed and is effective. 7.6.4.1.3. 7.6.4.1.4. Escalate the issue to a CAR (see Quality Manual). 7.6.5. Copies of QISs will be kept in maintenance, reagent, or standards binders, associated case packets, and/or electronically as appropriate. jnclu. 7.6.5.1. So included in case packets will be treated as notes.

8.0 PROCEDURES FOR COLOR TESTS

8.1 COLOR TESTS (General Procedure)

- Transfer small portions of the sample to the depressions of a white spot plate as needed.
- 8.1.2. Fransfer drop-wise volumes of each appropriate color test reagent(s).
- 8.1.3. Mix even(s) and sample when necessary.
- 8.1.4. Allow appropriate time to observe any color reaction. Document results in notes.
- 8.1.5. Run blanks when appropriate.
- 8.1.6 A standard may be ton for comparison.

8.2 COLOR TESTS (Specific Drug)

8.2.1. The results of color tests are presumptive and can direct subsequent confirmatory testing. The number of color tests used is at the discretion of the Criminalist and is dictated by the form of the substance. In general, a full set of color tests is not applied by the Criminalist. The color tests in bold are the ones typically used for color testing on suspected drug substances.

8.2.1.1. <u>Amine</u>	<u>s</u> Amphetamine:	Wagner: No Reaction Marquis: Orange or Crarge → Brown Nitroprusside: No Reaction Liebermann: Red → Drarge
	Methamphetamine:	Wagner: Brown Precipitate Marquis: Orange or Orange> prown Nitroprusside: Blue Liebermann: Orange
	MDMA:	Wagner: Brown precipitate Marquis: Grn and/or Purple —> Black Nitroprusside: Blue Mecke: Green/Blue

Approval: November 25, 2019



Ferric Chloride: No Reaction Duquenois/Chens #2: No Reaction Chens #2: No Reaction **CoSCN:** Blue +HCl: Light Green Lieberman: No Reaction

GBL:

Nor for LABORATORY USE

9.0 PROCEDURES FOR CRYSTAL TESTS

9.1 CRYSTAL TESTS (General Procedure)

Transfer a small portion of the sample to a slide or spot plate well as appropriate for analysis.

Add a small drop of the reagent(s) needed to produce crystals.

- 9.1.3 Mix the cample and reagent, if needed.
- 9.1.4 View with a microscope and record the results of the crystal formation.

9.2 CRYSTAL TEST'S (Specific Drug Procedures)

9.2.1. Amines

1.1

9.2.1.1 <u>Amphetamine/Manamphetamine</u>: Gold Chloride/Phosphoric Acid (Hanging Dr.p.C.ystal Test)

9.2.1.1.1.	Put a snel portion of the sample in a spot plate well
9.2.1.1.2.	Add a drop of saturated NaOH to the well and place a
	drop of gold where e hosphoric acid reagent on a
	microscope slide
9.2.1.1.3.	Invert the microscope slide over the spot plate well to
	allow the fumes produced from the sample to interact
	with the reagent on the slide
9.2.1.1.4.	Allow approximately 2 minutes for the fumes to react
	with the reagent (longer times may be necessary)
9.2.1.1.5.	Invert the slide and examine microscorically with a
	minimum of 100x magnification
9.2.1.1	.5.1. Amphetamine produces fan-shaped rystals (see
	photo on next page)
9.2.1.1	
	crystals (see photo on next page)
9.2.1.1	
	appear as clothespin crystals linked back to be ck
	(see photo on next page)



9.2.2 <u>Cocaine and Cocaine Base</u>: Gold Chloride (Direct)



9.2.4. <u>PCP</u>: Potassium Permanganate (KMnO₄)

- 9.2.4.1. Place a small amount of material on a microscope slide
- 9.2.4.2. Add a drop of 0.5N HCl or distilled water to the material
- 9.2.4.3. Add a few small $KMnO_4$ crystals to the slide and mix
- 9.2.4.4. Immediately examine microscopically with a minimum of 100x magnification

9.2.4.4.1. Crystals are pink bow-tie shapes (see photo below)



9.2.5 <u>GHB</u>: Silver Nitrate/Supric Nitrate

0 2 5 1	Dlace a small an	and the of material	on a microscope slide
9.2.5.1.	Place a Sillar al	our cor material	Sh a microscope shue

- 9.2.5.2. Add a drop of cease it to the material on the slide and mix
- 9.2.5.2.1. OR place a drop of the reagent near the sample and
 - combine via a "neck" by drawing one drop into the other
- 9.2.5.3. Crystals will often grow of the edges of the drop after about five minutes
- 9.2.5.4. Examine microscopically 9.2.5.4.1. Crystals are gray plate-life crystals which often are overlapping (see photo below)

GHB



10.0 IDENTIFICATION OF PHARMACEUTICAL PREPARATIONS

General Apparent Visual and Preliminary Identification Procedures

For a positive visual examination, the type of pharmaceutical (capsule/tablet), the color, shape, and logo/code must all be consistent with the reference description.

10.1.1.1. The following resources can be utilized for visual examinations of pharmaceutical preparations when the manufacturer's concellogo is clearly visible.

10.1.1.1.1. Drug ID Bible

10.1.1.1.2. **PX**-ID CD

10.1.1.1.3. Imprint code on the prescription label matching that isible in the tablet or capsule

- 10.1.1.1.4. On-live date identification websites
- 10.1.1.2. A photocopy of print at from a reference source is included in the case notes, when utilized for apparent tablet identifications.
- 10.1.1.3. If the visual examination read, to an apparent non-controlled medication, testing will be stopped at this point and the tablets/capsules will be listed and/or reported as containing an "apparent non-controlled medication"
- 10.1.1.4. If the visual examination leads to an apparent controlled substance, further testing is required in order to include the tablets/capsule in the report.

10.1.1.4.1. If the tablets/capsules are not going to be reported, no additional testing is required, but the notes must list the tablets or capsules as "apparent" controlled substance.

- 10.1.2. For sealed pharmaceutical preparations and liquids, including blister parks, the label information may be used for identification.
- 10.1.3. For the identification of pills without markings and liquids in unsealed containers, an instrumental or crystal test is required for identification.
- 10.1.4. If acetaminophen is identified in a tablet, the Criminalist must remove the acetaminophen, either chemically or by using an appropriate instrumental method, and retest the sample before reporting no controlled substance detected.

10.2 Final Identification Procedures

10.2.1 A final analysis requires a crystal test with an instrumental test, GC/MS with a standard, or GCMS with a second instrumental test.

Not for LABORATORY USE

11.0 GC/MS ANALYSIS

11.1 General Sample Preparation Guidelines



11.2 Specific Sample Preparations

11.2.1.	<u>Cocaine Base</u>		
	11.2.1.1.	Dissol	ve samples of cocaine base in hexane.
	11.2.1.2.	Analyz	ze using an appropriate method.
	1.05		
11.2.2.		.	
	11.2.2.1.	Liquid	
Λ.	11.2.2.	1.1.	Dissolve a small amount of sample into approximately 1 ml of methanol.
	11.2.2.	1.2.	Analyze using an appropriate method.
	11 2.2.2.	Sugar	cubes
	11.2.2.	2.1.	Examine under UV light (both short and long
			wavelength) to find the most concentrated spot of
			fluorescence
	1.2.2.	2.2.	Wash the concentrated area of the cube with methanol
			dropwise over the spot well.
	1 2.2.	2 3.	Collect the concentrated methanol and place into a test
			tube.
	11.2.2	2.4.	Dry down the concentrated methanol.
	11.2.2.	2.5	Reconstitute with a small amount of methanol.
	11.2.2.		Place the sample into a GCMS micro vial insert, and
			analyze using an appropriate method.
	11.2.2.3.	Blotte	r paper
	11.2.2.		Extract din city with a small amount of methanol.
	11.2.2.		Allow the sample to sit in the dark for at least 20
		-	minutes
	11.2.2.	3.3.	Place the sample is to a GCMS micro vial insert, and
			analyze using an appropriate method.
	11.2.2.4.	LSD w	ill be finalized due to the breakdown of the product over
		time.	
11.2.3.	<u>PCP</u>		
	11.2.3.1.	Plant	material,
	11.2.3.	1.1.	Extract by briefly placing a small appoint of plant material in a test tube and washing with methanol.
	11.2.3.	12	Vortex the sample for 5–10 seconds and allow the plant
	11.2.).	1.2.	material to settle.
	11.2.3.	12	Analyze using an appropriate method.
	11.2.		That yie doing an appropriate method.
	11.2.3.2.	Liquid	
	11.2.3.	-	Using a microcap, place one drop of the liquid into avia.
	11.2.3.		Add approximately 1.5-2 ml of methanol
	11.2.3.		Analyze using an appropriate method.
		,.	
11.2./	<u>Steroids</u>		
	11.2.4.1.	Using	a microcap, place one drop of the liquid into a vial.
	11.2.4.2.		pproximately 1 ml of methanol.
		A	

11.2.4.3. Analyze using an appropriate method.

- 11.2.4.4. To clean-up oil based steroids, an extraction can be done as follows:
 - 11.2.4.4.1. Vortex one part oil containing the steroid with two parts hexanes.
 - 11.2.4.4.2 If the oil and hexane mix, add the methanol to the oil/hexane mixture and vortex.
 - 11.2.4.4.2.1. The steroid will elute into the methanol.
 - 11.2.4.4.2.2. Pull off the methanol layer (top layer).
 - 11.2.4.4.2.3. Analyze using an appropriate method.
 - 11.2.4.4.3. If the oil and hexane solutions do not mix, the steroid will elute into the hexane.
 - 11.2.4.4.3.1. Pull off the hexane layer (hexane layer may be the upper or lower layer depending on the type of oil used in the preparation).
 - 11.2.4.4.3.2. Mix the hexane layer with methanol.
 - 1.2.4.4.3.3. The steroid will elute into the methanol layer.
 - **1.2.4.4.3.4**. Pull off the methanol layer (lower layer).
 - **2.** 4.3.5. Analyze using an appropriate method.
- 11.2.5. GHB/GBL/1,4-Bytanediol
 - 11.2.5.1. Place one drop of the liquid into a vial.
 - 11.2.5.2. Add approximately 1.5–2 ml of methanol.
 - 11.2.5.3. Analyze to p an appropriate method.
 - 11.2.5.4. Note: If GC/NS confirmation is positive for GBL and if GHB crystals were produced, results are reported as GHB. If no crystals were produced during the preliminary tests, results are reported as GHB/GBL.
- 11.2.6. <u>Tablets</u>
 - 11.2.6.1. Pulverize the tablet, or a portion of, into a fine powder (use enough of the tablet to get approximately 1mg of the active ingredient)
 - 11.2.6.2. Dissolve the material into approximately 1 rul of methanol in a test tube or vial.
 - 11.2.6.3. Mix the sample and then allow the material to settle.
 - 11.2.6.3.1. If the powdered sample was placed directly into a vial, the powder must not be more than approximately 0.25ml. If it is, the liquid must be transferred to a separate vial before instrumental analysis.
 - 11.2.6.4. Analyze using an appropriate method.
 - 11.2.6.5. To clean out unwanted fillers, utilize the following technique:
 - 11.2.6.5.1. Pulverize the tablet, or a portion of, into a fine powder.
 11.2.6.5.2. Make an aqueous solution by placing the powder into a test tube and adding approximately 2 ml of water.
 - 11.2.6.5.3. Make the aqueous solution basic by adding 1-2 drops of saturated NaOH.
 - 11.2.6.5.4. Vortex the solution and add approximately 1–2 ml of hexanes to the tube.

- 11.2.6.5.5. Vortex, then allow the two phases of the solution to separate.11.2.6.5.6. Remove the hexanes layer and place it into a GC/MS vial
 - for analysis.
- 11.2.6.5.7. Analyze using an appropriate method.

11.3 Methods

1.3.1. The current method parameters will be retained in the GCMS methods binder. The Criminalist will run the appropriate method for the suspected drug.

11.4 Reference Standards and Retention Time

- 11.4.1. Reference standards will be verified prior to being utilized for casework.
- 11.4.2. Reference spectra of drugs can be acquired and maintained on the GCMSs.
 - 11.4.2.1. Petertion time comparisons may only be used when the standard and questioned samples are run using the same instrument and method parameters.
 - 11.4.2.2. Retention times for standards are stable over a period of time, but must over established when the instrument conditions are significantly varied (change of method parameters, changing the column, e.c.).

11.5 GCMS Unknown Evaluations

- 11.5.1. Prior to comparisons to standards or libraries, the unknown data will be evaluated for suitability as follows:
 - 11.5.1.1. Negative controls do not include peaks of any drug or controlled substance. Any peaks in the blank are be related to column bleed, phthalates, etc.
 - 11.5.1.2. Unknown peaks of interest are single, smooth, symmetrical narrow peaks. Some tailing may be present.
 - 11.5.1.3.There is sufficient mass fragments for comparison and
identification without saturating the detector.
- 11.5.2. If the above criteria are not met, no comparisons will be conducted.
 - 11.5.2.1. Printouts do not need to be kept but the notes must reflect the testing that was conducted and why the results were not used for comparison.

- 11.5.3. Evaluations will be documented in the notes.
- 11.5.4. Samples may be reanalyzed to obtain better results (ex: different method, change in concentration).
- 11.5.5. The Criminalist will examine all integrated and significant nonintegrated peaks to determine if they are suitable for comparisons.

11.5.6. Any printouts not used to form final conclusions, interpretations, or opinions may be discarded, but the notes must indicate that the test was conducted and why the data was not kept.

n.o Reportion Time Identification

- 11.6.1. When the retention time of a unknown sample is being utilized as a confirmatory test, the following applies:
 - 11.6.1.1. The retention time of the standard was determined in-house.
 - 11.6.1.2. The clandard and the unknown were analyzed under the same instrumental conditions and method.
 - 11.6.1.3. The retention time of the unknown must be within \pm 5% of the retention unprof the reference standard.

11.7 Mass Spectra Identification

- 11.7.1. When comparisons are being made to the mass spectrum of an unknown, the following apply:
 - 11.7.1.1. Reference spectra acquired either on the instrument used or stored in a retrievable library (either computer or hard copy) may be used.
 - 11.7.1.1.1. An abbreviated/condensed abrar, spectrum should only be considered a tentative identification.
 - 11.7.1.2. The base peak and other prominent ions of the u known spectrum should match that of the reference.
 - 11.7.1.2.1. The relative intensities of the prominent ione should agree between the reference and the unknown spectrum.
 - 11.7.1.2.2. Prominent ions with a relative intensity greater than 10% of the base ion in the reference spectrum should be present (depending upon concentration) in the unknown spectrum.
 - 11.7.1.3. There should not be any major differences or additional prominent ions that are not explainable.
 - 11.7.1.4. The overall fragmentation pattern, and relative ion abundances are compared for consistency.

11.7.1.5. The mass spectrum of the unknown should contain the molecular ion, if present in the reference.

Nor cop LABORATORY US

Page 55 of 75

Approval: November 25, 2019

12.0 FTIR USER GUIDELINES

Analyzing Samples

Solids

12. 1.1.

12.1.1

12.1.1.3.

12.1

- Place enough sample to cover the sample window.
- Avoid damaging the sample window by minimizing its 12.1.1.1.1. contact with tools.
 - Appropriately close the Golden Gate attachment.
 - Run with an appropriate method.

Or the sample can be dissolved in an appropriate solvent prior 12.1.1.4. application onto the sample window and run as a liquid (see be ow).

12.1.2. Liquids

12.1.2.1.			ple on the sample window.
12.1.2.2.	Run with an a	ar prop	iate method.

12.1.3. Cast film from a volatile liquid sa nple

- 12.1.3.1.
- Place a drop of sample on the ample window. Allow to evaporate and leave a cast film on the sample window 12.1.3.2. (repeat to layer if necessary)
- Run with an appropriate method 12.1.3.3.
- 12.1.4. An appropriate background will be taken prior to each sample.
- 12.1.5. Unknown sample runs will be labeled with the incident of ase number, unique identifier, and the initials of the Criminalist working the case.
- 12.1.6. Instrument printouts are considered the original documentation
 - 12.1.6.1. Any printouts not used to form final conclusions, interpretations, or opinions may be discarded, but the pot must indicate that the test was conducted and why the data not kept.

12.2 FTIR Unknown Evaluations

- 12.2.1. Prior to comparisons to standards or libraries, the unknown data will be evaluated for suitability as follows:
 - 12.2.1.1. Blanks consisted mainly of broad peak complexes centered approximately around the areas of 2500 and 1300 cm⁻¹. They did not include unexpected significant peaks.
 - 12.2.1.2. The unknown spectra have smooth, well-formed peaks with appropriate reflectance percentages to allow for comparisons.
 - The fingerprint region contains a sufficient amount of peaks to allow for comparison.
- 12.2.2. If the rowe criteria are not met, no comparisons will be conducted.
 - 12.2.2.1. Printouts do not need to be kept but the notes must reflect the esting that was conducted and why the results were not used for comparison.
- 12.2.3. Evaluations will be documented in the notes.
- 12.2.4. Samples may be real-alyzed to obtain better results (ex: different change in concentration, different sample preparation).
- 12.2.5. Any printouts not used to form final conclusions, interpretations, or opinions may be discarded, but the notes must indicate that the test was conducted and why the data was not kept.

12.3 Library Matches

2.1.3.

- 12.3.1. When comparisons are being made to the FTIL spectrum of an unknown, the following apply:
 - 12.3.1.1. Reference spectra acquired either on the instrument used or stored in a retrievable library (either computer o) hard copy) may be used.
 - 12.3.1.1.1. An abbreviated/condensed library spectrum chould only be considered a tentative identification.
 - 12.3.1.2. The principle peaks of the unknown spectrum should nate that of the reference.
 - 12.3.1.2.1. The relative intensities of the principle peaks should agree between the reference and the unknown spectrum.
 - 12.3.1.3. There should not be any major differences or additional significant peaks that are not explainable.
 - 12.3.1.4. The peak pattern of the fingerprint region and relative intensities are compared for consistency.

12.4 Printing the Spectrum

12.4.1. At this time spectra should be printed in either black or dark blue, and at a line thickness of 3 or 4 to provide the best resolution when they are scanned for record retention. Requirements are subject to change.

12.5 Additional Information

12.5.1. The spectrum is recorded from 4000 – 650 cm ⁻¹, a minimum of 16 scans are used per sample, and the time the last background sample was run is automatically included on the FTIR printout.

13.0 Raman User Guidelines

13.1 Analyzing Samples

13.1.1. Solid or Liquid

1.1.2.

13.1.1.1. Place sample into an appropriate package, if not already in one.

Place the sample in its package over the circular opening on the universal platform sampling accessory ensuring the sample is over the window.

- Run an appropriate scan.
- 13.1.2. Unknown sample runs will be labeled with the incident or case number, unique identifier, and the initials of the Criminalist working the case.
- 13.1.3. Instrument printouts are considered the original documentation.
 - 13.1.3.1. Any printouts not used to form final conclusions, interpretations, or opinions may be discarded, but the notes must indicate that the test was conducted and why the data was not kept.

13.2 Raman Unknown Evaluations

- 13.2.1. Prior to comparisons to standards or libraries the unknown data will be evaluated for suitability as follows:
 - 13.2.1.1. The unknown spectra have smooth, well formed peaks with appropriate reflectance percentages to allow for comparisons.
 - 13.2.1.2. The unknown spectrum contains a sufficiency mount of peaks to allow for comparison
- 13.2.2. If the above criteria are not met, no comparisons will be conducted.
 - 13.2.2.1. Printouts do not need to be kept but the notes must reflect the testing that was conducted and why the results were not ased for comparison.
- 13.2.3. Evaluations will be documented in the notes.
- 13.2.4. Samples may be reanalyzed to obtain better results (ex: different packaging, different sample preparation).

13.2.5. Any printouts not used to form final conclusions, interpretations, or opinions may be discarded, but the notes must indicate that the test was conducted and why the data was not kept.

13.3 Library Matches

- 13.3.1. When comparisons are being made to the Raman spectrum of an unknown, the following apply:
 - 13.3.1.1.Reference spectra acquired either on the instrument used or
stored in a retrievable library (either computer or hard copy)
may be used.
 - 13.3.1.1.1An abbreviated/condensed library spectrum should only
be considered a tentative identification.
 - 13.3.2. The principle peaks of the unknown spectrum should match that of the reference.
 - The relative intensities of the principle peaks should agree between the reference and the unknown spectrum.
 - 13.3.1.3. There should not be any major differences or additional significant peaks that are not explainable.
 - 13.3.1.4. The overan peak pattern and relative intensities are compared for consistency

13.4 Printing the Spectrum

13.4.1 At this time, spectra should be printed in Einer black or dark blue to provide the best resolution when they are scanned for record retention. Requirements are subject to change.

13.5 Subtracting

- 13.5.1. When a sample to be analyzed is diluted by a known chernical, the chemical can be subtracted from the sample to identify the controlled substance present.
 - 13.5.1.1. Open the chemical spectrum, if saved, or add the spectrum to the same window as the diluted sample.
 - 13.5.1.2. Select both spectra in the display window and click on Subtra in the Process menu.
 - 13.5.1.3. The top spectrum should be the diluted sample. The middle spectrum should be the chemical to subtract. The bottom spectrum is the subtracted diluted sample
 - 13.5.1.3.1. If not, an arrow on the right of the window will switch the order.
 - 13.5.1.4.The subtracted spectrum is changed using the Factor Scale on
the left to subtract out the chemical to the desired level.

- 13.5.1.5. Once done, select "Add" to add the subtracted sample to a new window.
- 13.5.2. The sample spectrum can now be processed by following the Library Match and Printing steps.

Nor coplasson and we we have a set of the se

14.0 Botanicals

14.1 Cannabis/Concentrated Cannabis/THC

Cannabis cannot be confirmed at this time due to a percent THC requirement 14.1.1 in California law. All weights for suspected cannabis cases will be reported as gross weights due te California law. suspected cannabis cases will have only the active components reported once 14.1.3. confirmed (ex "contains THC" or "contains CBD"). 14.1.4. Analysis odified Duquenois-Levine Color Test 14.1.4.1. 14.1.4 ..1. Place some of the material into a test tube Cover the material with Duquenois reagent and mix 14.1.4.1.2 Add an approximately equal amount of concentrated HCl 14.1.4.1.3. and mix 14.1.4.1.3 A violet color should form Add an approximately equal amount of chloroform 14.1.4.1.4. A visitet color should form in the CHCl₃ layer 14.1.4.1.3.1. This test call also be run by first extracting the material 14.1.4.1.5. with petroleum ether, which is then dried down prior to the addition of the Inquenois reagent. GCMS 14.1.4.2. A small portion of the sample can be extracted with 14.1.4.2.1. methanol and analyzed with an appropriate GC/MS method. Or, a small portion of the sample can be extracted with 14.1.4.2.2. petroleum ether, dried down, reconstruted with methanol, and analyzed with an appropriate GC/MS method. C'SA **Extraction of Cannabinoids from Food Products** 14.1.4.3. **Reagents**: 14.1.4.3.1. 0.2N Methanolic KOH 1.0 N HCl 10% Ethyl Acetate: Hexane **Extraction Procedure:** 14.1.4.3.2. Add sample to a 15ml screw-cap conical vial. 14.1.4.3.2.1. Add approximately 4ml hexane to sample vial and 14.1.4.3.2.2. vortex until the sample is dissolved. Add approximately 4ml hexane to an empty vial 14.1.4.3.2.3. to act as a negative control.

- 14.1.4.3.2.4. Add approximately 4ml 0.2N methanolic KOH to each vial and shake for 5 minutes.
- 14.1.4.3.2.5. Centrifuge to separate the layers.
- 14.1.4.3.2.6. Remove and discard the top hexane layer.
- 14.1.4.3.2.7. Add approximately 6ml hexane to each vial and shake for 5 minutes.
- 14.1.4.3.2.8. Centrifuge and discard the top hexane layer.
- 14.1.4.3.2.9. Repeat steps 14.1.4.3.2.7. and 14.1.4.3.2.8.
- 14.1.4.3.2.10. Add approximately 1ml deionized water to each vial.
- 14.1.4.3.2.11. Add approximately 6ml hexane to each vial and shake for 5 minutes.
- 14.1.4.3.2.12. Centrifuge and discard top hexane layer.
- 14.1.4.3.2.13. Add approximately 1.5ml 1.0N HCl to each vial and check pH to ensure the solution is acidic. Add more HCl if necessary.
- 14.1.4.3.2.14. Add approximately 3ml 10% ethyl acetate: hexane to each vial and shake for 5 minutes.
 - 4.1.4.3.2.15. Centrifuge to separate layers.
 - .1. 3.2.16. Transfer top organic layer to a new tube.
- 14 3.2.17. Dry down the organic layer and reconstitute with methanol.

14.1.4.2 2.18. Analyze using an appropriate GCMS method.

<u>Reference</u>: AFE, Department of Defense Drug Detection Quality Assurance Laboratory TNC Quantitation in Hemp Oil.

14.2 Psilocin/Psilocybin

- 14.2.1. A preliminary result for psilocin/psilocyble consists of positive results for both a color test and microscopic examination of the pores, or a positive instrumental test.
- 14.2.2. A final result for psilocin/psilocybin consists of positive preliminary results and an additional positive instrumental result.
- 14.2.3. Analysis

10,

- 14.2.3.1. Macroscopic Characteristics Common to Psilocin/Delocybin Mushrooms
 - 14.2.3.1.1. Fluted stem
 - 14.2.3.1.2. Inky blue coloring on various areas of the stem
 - 14.2.3.1.3. Gold colored crinkled cap
 - 14.2.3.1.4. Dark gills

S



14.2.3.4.1.2. For fresh plant material, it may be necessary to place a mushroom cap in a plastic cup containing

- liquid nitrogen and allow the mushroom to equilibrate prior to grinding.
 14.2.3.4.2. Soak in methanol for at least one hour.
 14.2.3.4.3. Analyze using the appropriate method.
- 14.2.3.4.4. Mushroom samples negative for psilocin/psilocybin must be soaked in methanol overnight prior to GCMS analysis in order to be reported as NCSD.

- 14.3.1. The presence of cathinone, with or without cathine/phenylpropanolamine, indicates that the plant material is Khat.
- 14.3.2. A preliminary result consists of a positive result for the presence of cathinone and/or cathine/phenylpropanolamine with GCMS analysis.
- 14.3.3. A final result consists of the preliminary result and a GCMS retention time comparison of the upknown and the appropriate standards of cathinone and/or cathine/phenylpropapolamine.
- 14.3.4. If only cathine/phenylprop notamine is found, the report will reflect those substances and will not differentiate between the two.
- 14.3.5. Extraction and GCMS analysis
 - 14.3.5.1. Use approximately 4 grap s ordried material or chop up fresh material to obtain the same amount.
 - 14.3.5.2. Place material in a 250 ml arlend ever flask and cover with 0.2N H_2SO_4 .
 - 14.3.5.3. Add approximately the same an ount of $0.2N H_2SO_4$ to a second flask as a negative control.
 - 14.3.5.4. Sonicate flasks in water bath for 30 m nutes
 - 14.3.5.5. Pour off liquid through filter funnel into a second 250 ml Erlenmever flask
 - 14.3.5.6. Make solutions basic with concentrated NaOH (dro wise until a color change is noted).
 - 14.3.5.7. Add approximately 30 ml of chloroform to each Mask. Mix well.
 - 14.3.5.8. Remove the chloroform layer (bottom) using separatory lunnel or a by pipetting into large test tubes.
 - 14.3.5.9. Evaporate to dryness using air.
 - 14.3.5.10. Reconstitute in no more than 2 ml of methanol.
 - 14.3.5.11. Analyze using an appropriate GCMS method.

14.4 OPIUM/OPIUM POPPIES

14.4.1. Extraction Procedure for Poppies

14.4.1.1. <u>Fresh Poppies</u>:

^{14.2.3.5.} When identification is made using GCMS, the results are reported as "psilocin/psilocybin."

- 14.4.1.1.1. Score and extract the sap from the pods
- 14.4.1.1.2. Dissolve the sap in methanol.
 - 14.4.1.1.2.1. Or, freeze dry the pods with liquid nitrogen and grind them into a powder.
 - 14.4.1.1.2.2. Extract powder in methanol about 20 minutes.

PLADORATORY USE

- 14.4.1.1.3. Analyze using an appropriate GCMS method.
- 14.4.1.2. <u>Dried Poppies</u>:
 - 14.4.1.2.1. Grind pods to a powder.
 - 14.4.1.2.2. Extract powder in methanol about 20 minutes.
 - 14.4.1.2.3. Analyze using an appropriate GCMS method.

.4.2 Three of the five principal alkaloids found in opium (morphine, codeine, the naine, noscapine, and papaverine) must be present before the sample can be classified as containing opium, or the poppies can be classified as opium poppier, *Papaver somniferium*.

15.0 COURT

15.1 GENERAL

15.1.1. General court policies are covered by the following references:

Λ	15.1.1.1.
	15.1.1.2.
- 🖊	15.1.1.3.

- Quality Manual City of San Diego Employee Code of Conduct Handbook
- SDPD Procedure 1.11

15.2 **TESTIMONY REGARDING EFFECTS**

- 15.2.1. Testimony to the physiological effects of seized drug substances are handled by the op-call detective experts in the narcotics section or Bio-Tox.
- 15.2.2. Criminalists call only testify about general effects of classes of drugs.

15.3 COURT EVALUATIONS

- 15.3.1. Evaluations will be done a minimum of once per accreditation cycle in each discipline.
- 15.3.2. Evaluations will be performed by mother qualified Criminalist.
- 15.3.3. If a criminalist has not testified in a discipline during the accreditation cycle, they will notify the QA Manager by email.
- 15.3.4. Evaluation forms or emails are kept by the QA Janager.

15.4 COURT POLICY

- 15.4.1. Criminalists generally operate on an "on-call" basis and should not appear on the basis of a subpoena alone.
- 15.4.2. A criminalist should be placed on-call by the subpoenaing agency when the actual date of the trial is finalized and no later than the day before they are needed to allow time to prepare the court packet.
- 15.4.3. The subpoenaing agency should maintain close communication with the Criminalist on the day needed and allow a one-hour response time for court.
- 15.4.4. If a Criminalist is unavailable for court, the unit supervisor will assign another Criminalist to reanalyze the case or have the technical reviewer testify.
- 15.4.5. When a Criminalist is planning to be away from the office for three or more business days, they must have an out of office memo issued to the district and

city attorneys, put an out of office autoreply on their email, and change their voicemail to an out of office message for the duration of their absence.

15.5 **PROCESSING SUBPOENAS FOR DRUG CASES**

15.5.1. Subpoenas arrive in batches and are logged in by trial date. The clerical staff processes and places them in the Forensic Chemistry bin for dissemination as needed.

Each Criminalist is responsible to follow-up on their subpoenas.

- The Criminalist will need to track their cases electronically and 15.2.1. ensure final reports are completed prior to set trial dates.
- 15.5.3. Court cases can be tracked as follows:
 - Frial-Jury court cases can be checked for readiness hearing 15.5.3.1. lates in SDLaw.
 - To 15.5.3.2. cess SDLaw, go to the department intranet. Enter your ID# and your password then select "Log In".
 - Type in DA10 followed by a space and then the 15.5.3.2.1. Prosecutor's Case Number listed on the subpoena and hit einer
 - If the Pres. No. has an "M" or no letter, a Q must be 15.5.3.2.2. added o the front.
 - The number must be 7 digits long. If only 5 are listed, 15.5.3.2.3. add a 01 (1st defendant), 02 (2nd defendant), etc, at the end.

15.5.3.3.

- Write the readiness date or leach subpoena to follow up.
 If no readiness hearing to listed, you must assume the case is "set"/"confirmed" unless you find out otherwise. 15.5.3.3.1. Sometimes the readiness will have taken place before 15.5.3.3.2. you check it on the computer. If so, he results will be listed.
- All "Superior Court" or "Jury-Trial" cases car be checked on 15.5.3.4. the computer a couple of days following the readinger. If not listed, the appropriate phone number on the subpoena must be contacted.
- Subpoenas with the following readiness result do not need to 15.5.3.5. finalized and should be marked as such and filed:

VACATED PC1000 CALLED OFF PLEAD GUILTY PRELIMINARY EXAM CONTINUED SENTENCING INFORMATION OTHER DISCRETIONARY REASON

ERROR CHANGE EVENT

15.5.3.6. Any subpoenas with the following readiness results must be finalized prior to the trial date:

SET Nor top LABORATORY USE CONFIRMED

Page 69 of 75

Approval: November 25, 2019

16.0 REAGENT PREPARATION

16.1.	<u>0.2N H₂SO4</u>	
	16.1.1.	Add 2.8ml concentrated H ₂ SO ₄ to approximately 300ml of DI H ₂ O
		in a volumetric flask.
Λ	16.1.2.	Bring up to volume with DI H ₂ O.
	0.2N Methan	
1 .2.	16.2.1.	Dissolve 5.611g KOH into a 500ml flask containing
	11.2.1.	approximately 400ml of MeOH.
	10.2.2.	Bring up to volume with MeOH.
•		
16.3.	<u>1.0NAC</u>	
	16.3.1.	Add 42ml concentrated HCl to 500ml flask containing
		pproximately 300ml of DI H_2O .
	16.3.2.	Pany up to volume with DI H ₂ O.
16.4.	<u>0.5N HCl</u>	
10.4.	<u>16.4.1.</u>	Add 21rd concentrated HCl to 500ml flask containing
	10.4.1.	approximately 300ml of DI H_2O .
	16.4.2.	Bring up to volume with DI H_2O .
16.5.	<u>10% Ethyl Ac</u>	<u>etate: Hexane</u>
	16.5.1.	Combine 10ml etbyl accepte with 90ml hexane.
16.6.	<u>Chens 2 (1%)</u> 16.6.1.	Dissolve 1 gram of cupric suffice in 100ml of H ₂ O.
	10.0.1.	Dissolve I grain of cupic storte in room of H ₂ O.
16.7.	Cobaltous Th	iocyanate (CoSCN)
	16.7.1.	Dissolve 1.0 grams of cobalt ace ate a to 1.5 grams potassium
		thiocyanate in 90ml H ₂ O.
	16.7.2.	Add10 ml of glacial acetic acid.
	16.7.3.	Add 100ml of glycerin.
	16.7.4.	Mix thoroughly
46.0	Duquanaia I	
16.8.	<u>Duquenois-L</u> 16.8.1.	
	10.8.1.	Dissolve 2.8 grams Vanillin and 5.8ml Acetaldehyde in 200ml of 200 proof Ethanol.
	16.8.2.	Bring total volume to 400ml with additional ethanol.
	10.0.2.	bring total volume to 400mm with adultional ethanol.
16.9.	<u>Ferric Chloric</u>	de The second
-	16.9.1.	Dissolve 10.0 grams of ferric chloride in 100ml of H ₂ O.
16.10.	Gold Chloride	
	16.10.1.	Dissolve 1.0 grams of gold chloride in 20ml of H_2O

1

Page 71	of 75	Seized Drug Manual Printed copies are not controlled	Approval: November 25, 2019
	16.21.2.	This solution must be made the day of	use.
	16.21.1.	Dissolve a small amount of Fast Blue E H_2O . Color should be light straw.	3 in approximately 2ml of
16.21.	<u>Weber</u>		
	16.20.2.	in 5ml of DI H ₂ O. Bring volume to 100ml with DI H ₂ O.	
10.20.	<u>Wagner</u> 16.20.1.	Dissolve 1.27 grams of iodine and 2.75	grams potassium iodiac
16 20	<u>Wagner</u>	no proparation, rotacoram pormangar	
16.19.	<u>Potassium Pe</u> 16.19.1.	<u>rmanganate (KMnO4)</u> No preparation, Potassium permangar	nate used in crystal form
16.18.	<u>Platinum (Pla</u> 16.18.1.	<u>itinic) Chloride</u> Dissolve 1.0 gram of chloroplatinic acio	d in 20ml of 4. 0.
	16.17.2	2.1. Dissolve 2.0 grams of sodium d	rbonate in 100ml of H₂O.
	16.17.2.	Prepare 2% sodium carbonate	
	16.17.1 16.17.1		
	16.17.1	.1. Dissolve 2.0 grams of sodium n MeOH.	itroprusside in 40ml of
10.17.	16.17.1.	Prepare sodium nitroprussi lo	
16 17	<u>Nitroprusside</u>		
	16.16.3.	Let sit overnight, there should be a thi the bottom.	n crystal layer settled on
	16.16.2.	Continue as ding approximately 5g of 1 crystals no longer dissolve completely.	
	16.16.1.	Add approximately 5g of NaOH solid to	
16.16.	NaOH (Satura	ited)	
	10.15.1. 16.15.2.	Ad 7ml of concentrated HCl.	/3111 01 1120.
16.15.	<u>Mercuric Iodi</u> 16.15.1.	Desolve 5 grams of Mercuric Iodide in t	72ml of H2O
		sulfuric acid.	
16.14.	<u>Mecke</u> 16.14.1	Dissolve 0.25 grams of selenious acid i	n 25ml of concentrated
	` \	and 40% Formaldehyde kept in separa	te containers.
10.13.	<u>Marquis</u> 1616.13.1.	No preparation. Marquis reagent is con	ncentrated Sulfuric Acid
Λ.		concentrated sulfuric acid.	
16.12.	<u>Lieberman's</u> 16.12.1.	Dissolve 10.0 grams of potassium nitri	te in 100ml of
		acid.	I
	16.11.1 16.11.2.	1. Add 13.2ml of conc. phosphoric Dissolve 1.0 grams of gold chloride in 2	
16.11.	<u>Gold Chloride</u> 16.11.1.	<u>/phosphoric Acid</u> Prepare 1+2 phosphoric acid	
16 11	Cold Chlorida	haspharic Acid	

17.0 APPROVED ABBREVIATIONS

Definition	Abbreviation (no regard to capilalization or periods)
side) and side two (tab/cap)	<>
Acetammophen	APAP, ACETA
Administrative Review	A, AR
Alprazolam	Alp
Aluminum	Al
Amount	Amt
Apparent	Арр
Barcode	BC
Black	Blk
Blue	Bl
Bottle	Btl
Brown	Pn
Cannabidiol	
Capsule, Capsules	Car, Cars
Carry over	с/о
Cellophane	Cello
Change due to tech review	TR
Chunky material	СНМ
Chunky white material	СѠМ
Chunky powder	СНР
Chunky powder material	СНРМ
Cigarette	Cig
Cocaine	Сос
Combined	Comb.
Concentration	Conc
Container	©

Container containing	C ²
Containing	c
crystal/crystalline material	СМ
Cut to open	СТО
Did not label	DNL
Each	ea
Emelope	Env
Estinlated	Est
Faint	Ft
Federally	Fed
Federally Contrained	FC
Federally Controlled Medication	FCM
Flash	Fl.
Fragment	Frag
Factory Sealed Packet	FSP
Green	Grn or Gr
Gross weight	GW
Heat Sealed	HS
Heroin	Ler
Hexane	The second se
Hydrocodone	Hydro
Initial Color Observed, Turned to Second Color Color containing a color, or 2 sides of tablets or capsules	
Identification	ID
Including/ Included	Inc.
knotted	k
Knotted plastic	kp
Knotted plastic baggie(s)	Kpb(s)
Labelled to contain	LTC
Light	Lt.
Manila Envelope	ME
Marijuana	MJ

Material	Mtl, Mtrl, Matl, Mat
Methamphetamine	Meth
Net weight	NW
No Change	Νο Δ
No Controlled Substance Detected	NCSD
No initials on seal	NIOS
No Reaction	NR, No Rx, -
Not 7/non Controlled	NC
Not altion Controlled Substance	NCS
Non-Controled Medication	NCM
Not Laboratory Lxa. nined	NLE
Orange	Or, O, and Org
Over the Counter	OTC
Oxycodone	Оху
Paraphernalia	Para
Piece	рс
Plastic Baggie	РВ
Positive Reaction	
Paper	pr
Paper bag	TN
Plant	Plut
Powder material	РМ
Precipitate	Ppt
Prescription	Rx
Prosecution	Pros
Purple	Purp
Reaction	Rxn
Rectangular	Rec.
Residue	Res
Residue amount of debris	RAD
Rock-like material	RM
Room Temperature	RT

Sealed	Sld
Slight	Slt.
Small	Sm
Tablet, Tablets	Tab, Tabs
Tar/Tar-like material	TM
Technical Review	Т
Tetal Gross Weight	TGW
Total Not Weight	TNW
Very	V
Volume	Vol
Weight	Wt
Ziploc	Zip
Ziploc Plastic Baggre	Zip Pb, zpb

Approval: November 25, 2019